

COUNTDOWN TO 2015 DECADE REPORT (2000–2010)

with country profiles

Taking stock of maternal, newborn and child survival

WHO Library Cataloguing-in-Publication Data

 $Countdown \ to \ 2015 \ decade \ report \ (2000-2010): \ taking \ stock \ of \ maternal, \ newborn \ and \ child \ survival.$

1. Women's health. 2. Child welfare - trends. 3. Infant welfare - trends. 4. Infant, Newborn. 5. Child. 6. Maternal mortality. 7. Infant mortality. 8. Survival. 9. Health priorities. 10. Millennium development goals. I. World Health Organization. II. UNICEF.

ISBN 978 92 4 159957 3 (NLM classification: WA 310)

© World Health Organization and UNICEF 2010

All rights reserved. Publications of the World Health Organization can be obtained from WHO Press, World Health Organization, 20 Avenue Appia, 1211 Geneva 27, Switzerland (tel: +41 22 791 3264; fax: +41 22 791 4857; email: bookorders@who.int).

The World Health Organization and UNICEF welcome requests for permission to reproduce or translate their publications — whether for sale or for noncommercial distribution. Applications and enquiries should be addressed to WHO, Office of Publications, at the above address (fax: +41 22 791 4806; email: permissions@who.int or to UNICEF, 3 United Nations Plaza, New York, NY 10017, USA.

The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the World Health Organization or UNICEF concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by the World Health Organization or UNICEF in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

The World Health Organization and UNICEF do not warrant that the information contained in this publication is complete and correct and shall not be liable for any damages incurred as a result of its use.

Printed in Washington, DC.

Cover photo, WHO/Mark Kokic; page iii, UNICEF/NYHQ2007-2758/Claudio Versiani; page 3, Save the Children/Joshua Roberts; page 4, UNFPA/Peter Bruyneel; page 9, UNFPA/Don Hinrichsen; page 12, UNICEF/NYHQ2006-2860/Julie Pudlowski; page 14, UNICEF/NYHQ2005-2402/Anita Khemka; page 16, UNFPA; page 18, UNICEF/NYHQ1992-0311/Sean Sprague; page 33, UNFPA/Ken Opprann; page 35, Photoshare/Rezaul Haque; page 39, Save the Children/Jonathan Hubschman; page 40, UNICEF/NYHQ2005-2410/Anita Khemka.

Editing and layout by Communications Development Incorporated, Washington, DC.



COUNTDOWN TO 2015 DECADE REPORT (2000–2010)

with country profiles

Taking stock of maternal, newborn and child survival



Contributors

Production editor: Jennifer Requejo (PMNCH/Johns Hopkins University)

Lead writers: Jennifer Requejo (PMNCH/Johns Hopkins University), Jennifer Bryce (Johns Hopkins University)

Subeditors/writers: Joy Lawn (Save the Children/Saving Newborn Lives), Peter Berman (World Bank), Bernadette Daelmans (WHO), Laura Laski (UNFPA), Cesar Victora (Universidade de Pelotas), Elizabeth Mason (WHO)

Profile support team: Tessa Wardlaw (UNICEF), Holly Newby (UNICEF), Archana Dwivedi (UNICEF)

Additional writing team: Henrik Axelson (PMNCH), Zulfiqar Bhutta (Aga Khan University), Ties Boerma (WHO), Flavia Bustreo (PMNCH), Eleanora Cavagnero (WHO), Mickey Chopra (UNICEF), Giorgio Cometto (Global Health Workforce Alliance), Andres de Francisco (PMNCH), Helga Fogstad (Norad), Giulia Greco (LSHTM), Kate Kerber (Save the Children/Saving Newborn Lives), Blerta Maliqi (WHO), Tim Powell- Jackson (LSHTM), Lale Say (WHO), Ann Starrs (Family Care International), Nancy Terreri (FCI/PMNCH)

Countdown Coordinating Committee

Mickey Chopra (co-chair), Zulfiqar Bhutta (co-chair), Jennifer Bryce, Joy Lawn, Tessa Wardlaw, Elizabeth Mason, Bernadette Daelmans, Helga Fogstad, Flavia Bustreo, Ann Starrs, Laura Laski, Ties Boerma, Cesar Victora, Peter Berman, Andres de Francisco

Technical working groups

Coverage: Jennifer Bryce (co-chair), Tessa Wardlaw (co-chair), Holly Newby, Archana Dwivedi, Jennifer Requejo, Alison Moran, Shams El Arifeen, Edilberto Loiaza, Angella Mtimumi, Blerta Maliqi, Lale Say, James Tibenderana, Anuli Ajene

Equity: Cesar Victora (co-chair), Ties Boerma (co-chair), Aluisio Barros, Carine Ronsmans, Wendy Graham, Edilberto Loiaza, Betty Kirkwood, Zulfiqar Bhutta, Kate Kerber, Henrik Axelson

Financing: Peter Berman (chair), Henrik Axelson, Giulia Greco, Ruth Levine, David Collins, Giorgio Cometto, Jacqueline Mahon, Ravi Rannan-Eliya, Josh Lozman, Nouria Brikci, Karin Stenberg, Dan Kraushaar

Health systems and policies: Bernadette Daelmans (co-chair), Helga Fogstad (co-chair), Yves Bergevin, Geoff Black, Nouria Brikci, Craig Burgess, Naomi Cassirer, Eleonora Cavagnero, Mickey Chopra, Giorgio Cometto, Vincent Fauveau, Lynn Freedman, Neeru Gupta, Julia Hussein, Monir Islam, Dan Kraushaar, Blerta Maliqi, Elizabeth Mason, Zoe Matthews, Barbara McPake, Robert Scherpbier, Anuraj Shankar, Nancy Terreri, Mark Young, David Sanders, Yaron Wolman

Acknowledgements

The *Countdown* group would like to thank the following:

UNICEF/Statistics and Monitoring Section for use of global databases, preparation of country profiles and review of report text. Particular recognition goes to Priscilla Akwara, David Brown, Danielle Burke, Rouslin Karimov, Rolf Lugendijk and Danzhen You for their input and review of the report.

The Countdown advocacy subcommittee chaired by the PMNCH and Family Care International for their inputs in shaping the headlines, action points, conclusion and media strategy: Flavia Bustreo (co-chair), Ann Starrs (co-chair), Adam Deixel, Debra Jones, Jacqueline Toupin, Lori McDougall, Nancy Terreri and Jennifer Requejo.

The PMNCH secretariat for convening meetings and teleconferences for the Countdown and PMNCH colleague Dina el Husseiny for providing administrative support.

UNFPA colleagues Stan Bernstein, Howard Friedman and Hedia Belhadj for their inputs and review of the report text.

The Lancet for preparation of the Countdown map and permission to reproduce the cause of child death pie chart, 2008.

Julia David at Johns Hopkins University for her work on the initial analysis of the *Countdown* coverage databases.

AusAID, Gates Foundation, Norad, SIDA, DfID, World Bank and USAID for providing financial support.

Christopher Trott and Elaine Wilson from Communications Development Incorporated for their work on copyediting and typesetting the report.

LSHTM for hosting the Core Group meeting in 2009. And all Countdown Core Group members: Andres de Francisco, Andy Haines, Ann Starrs, Anne Mills, Anurai Shankar, Archana Dwivedi, Anne Tinker, Anthony Costello, Bernadette Daelmans, Betsy McCallon, Betty Kirkwood, Blerta Maligi, Carine Ronsmans, Ties Boerma, Cesar Victora, Daniel J. Carucci, David Sanders, Debra Jones, Donna Bowers, Doyin Oluwole, Edilberto Loaiza, Elizabeth Mason, Flavia Bustreo, Frank Smith, Giorgio Cometto, Hamid Rushwan, Hedia Belhadi, Helga Fogstad, Henrik Axelson, Jaime Sepulveda, Jim Tulloch, Jeffrey Sturchio, Jennifer Bryce, Jennifer Requejo, Joanna Schellenberg, Jo Nicholls, Josh Lozman, Joy Lawn, Julia Hussein, Kate Eardley, Katie Porter, Lale Say, Lars Gronseth, Leora Hanser, Lori McDougall, Margaret Chirgwin, Maria Francisco, Mary Bassett, Marta Seoane Aguilo, Monir Islam, Michael Klosson, Micky Chopra, Nancy Terreri, Neil Pakenham-Walsh, Oona Campbell, Peter Berman, Regina Keith, Richard Horton, Rifat Atun, Sadia Chowdhury, Simon Wright, Steven Hodgins, Stan Bernstein, Stewart Tyson, Tessa Wardlaw, Tim Shorten, Timothy Powell-Jackson, Vincent Fauveau, Wendy J. Graham, Zoe Matthews, Zulfigar Bhutta.



Countdown to 2015: key messages for 2010

- The Countdown report for 2010 contains good news—many countries are making progress, reducing mortality and increasing coverage of effective health interventions at an accelerating pace.
- But the news is not all good. Many Countdown countries are still off track for achieving Millennium Development Goals 4 (reduce child mortality) and 5 (improve maternal health), and are not increasing coverage of key health interventions guickly enough.
- Countdown countries in Sub-Saharan Africa are especially far behind, although a few have shown improvements.
- The vast majority of maternal and child deaths are preventable, but unacceptably large numbers of women, newborns and children are still dying each year in *Countdown* countries, where at least 95% of all maternal and child deaths occur. A growing proportion of child deaths occur in the first four weeks of life.
- Poorly functioning health infrastructure, inadequate numbers of health workers, slow adoption of evidence-based health policies and insufficient focus on quality of care are holding back progress in many countries.

- Skilled care at birth, including emergency care for mothers and newborns, is critical to achieving Millennium Development Goals 4 and 5: about 2 million lives a year are lost to complications occurring during labour and childbirth.
- Pneumonia and diarrhoea remain the largest killers of children after the newborn period.
 Undernutrition contributes to more than onethird of child deaths.
- Some Countdown countries are doing better at reaching the most disadvantaged women and children, but profound inequities in coverage and health outcomes—both between and within countries—must be confronted and overcome.
- Countries should aggressively pursue policies to make health services available and affordable for all, by making services free at the point of delivery and exploring innovative financing strategies.
- Funding is increasing for maternal and child health, but at too slow a pace, and funding for family planning has declined.
- Millennium Development Goals 4 and 5 are still achievable by 2015—but only a dramatic acceleration of political commitment and financial investment can make it happen.





35

Contents

Countdown headlines for 2010: saving the lives of the world's women, newborns and children

About Countdown to 2015 5

Progress towards Millennium Development Goals 4 and 5 **7**

Causes of maternal and child deaths 11

Social determinants of maternal, newborn and child health **13**

Coverage along the continuum of care 15

The building block framework: health systems and policies to save the lives of women, newborns and children 28

Equal care for every mother and child 31

Country spotlight:

Brazil's success in narrowing the gap

Country spotlight:

Narrowing gender differentials in Bangladesh

Financing for maternal, newborn and

child health 36

Action now for Millennium Development

Goals 4 and 5 **38**

Conclusion 40

Country profiles 41

Notes **191**

References 192

Abbreviations

BCG Bacille Calmette-Guérin

DPT diphtheria and tetanus with pertussis

GDP gross domestic product

HIV/AIDS human immunodeficiency virus/acquired immune deficiency syndrome

MDG Millennium Development GoalODA official development assistance

UNAIDS Joint United Nations Programme on HIV/AIDS

UNIFPA United Nations Population Fund UNICEF United Nations Children's Fund World Health Organization

Countdown headlines for 2010: saving the lives of the world's women, newborns and children



Survival status

Millennium Development Goal 4—reduce child mortality

- Good news: 19 of the 68 Countdown countries are on track to achieve Millennium Development Goal (MDG) 4.
 - 17 countries have reduced child mortality by at least half.
 - 47 countries have accelerated their progress on child mortality since 2000.
- Much work remains: 49 Countdown countries are not on track to achieve MDG4.
 - 12 countries (including some currently on track) have seen their progress slow since 2000.
- Death and illness:
 - Globally 8.8 million children a year die before their fifth birthday, more than 40% of them during their first four weeks of life. At least two-thirds of all child deaths are preventable.
 - Pneumonia and diarrhoea remain the largest killers of children after the newborn period.
 - Undernutrition contributes to more than 1 in 3 child deaths.

Millennium Development Goal 5—improve maternal health

- Good news: new studies suggest that some progress is being made on reducing maternal mortality.
- Much work remains: both globally and in most Countdown countries, progress is insufficient to achieve MDG 5, particularly in Sub-Saharan Africa. Urgent action is needed to scale up proven interventions to improve reproductive and maternal health.

Death and illness:

- An unacceptable number of women die in pregnancy and childbirth each year.
 For every woman who dies, at least 20 others suffer injuries, infection and disability. Almost all maternal deaths are preventable.
- Most maternal deaths occur during childbirth and in the immediate postnatal period, which is also when most stillbirths and newborn deaths occur.
- The leading cause of maternal deaths remains postpartum haemorrhage, largely preventable through skilled care during childbirth.

Coverage gains and gaps

- Progress is inconsistent: progress on coverage of lifesaving interventions across the continuum of care is uneven.
 - Some interventions delivered routinely through outreach or scheduled in advance (such as vaccinations and vitamin A supplementation) have achieved and sustained high coverage.
 - Interventions that must be provided in response to acute need (such as treatment of childhood illnesses and caesarean sections) show little progress.
 - Relatively new interventions that have received attention and resources, such as insecticide-treated nets and prevention of mother-to-child transmission of HIV, show rapid gains.
- Skilled care during childbirth: all women and newborns need access to a skilled attendant at birth, but overall coverage across the *Countdown* countries remains insufficient and uneven.
 - 10 countries showed coverage gains of more than 10 percentage points since 1990, and 3 countries—Burkina Faso,

- Pakistan and Rwanda—had gains of more than 20 percentage points from around 2000 to around 2008.
- 11 countries have shown no progress in coverage since 1990.
- Family planning: wide disparities in coverage of family planning services across and within countries represent a missed opportunity to improve the health of women and young children.
- More information is needed: higher coverage is critical, but saving lives also depends on the quality of care. More information is needed on what care is actually provided during antenatal, childbirth and postnatal contacts.

Health systems and policies

- Health workers: 53 of the 68 Countdown
 countries are experiencing acute shortages
 of doctors, nurses and midwives.
 Overcoming these shortages and
 addressing the unequal distribution of
 health workers within countries require
 focused investment in training, deployment
 and retention.
- Financial barriers to access: the high proportion of health service costs paid out of pocket in nearly all *Countdown* countries puts families at risk of financial catastrophe. Making services free at the point of delivery helps increase utilization: financing mechanisms such as prepayment and risk pooling can help make health services available and affordable for all.
- Improving access and quality of care: investment in health information and referral systems, equipment, medical supplies and infrastructure is critical to improving access to and quality of maternal, newborn and child health services.
- Adoption of policies: evidence-based policies can save and improve women's and children's lives. Bangladesh and Nepal, for example, have shown that implementing policies to increase access to diarrhoea and pneumonia treatment in the community reduces child deaths. More progress is needed: the number of Countdown countries that have adopted

recommended policies for increasing access to quality care is still too low.

Closing the equity gap

- Inequities in access: coverage rates are substantially higher among women and children in better-off families than in poor families.
- The poor and excluded: high national coverage levels do not always indicate progress in reaching the poorest and most vulnerable women and children. Guatemala and Zambia, for example, have similar levels of overall coverage for a subset of proven maternal, newborn and child health interventions, but more women and children from the poorest families receive these services in Zambia than in Guatemala.
- Further research needed: countries with smaller gaps between rich and poor including Bangladesh, Brazil, Egypt, Swaziland and Zambia—may provide models for reducing inequities through greater political commitment, specific targeting of low-income groups, redirecting of human resources and other strategies.
- Service provision: disparities are larger for services provided in health facilities (such as delivery care) than for those delivered at the community level (such as vaccines).

Closing the funding gap

- Financing the gap: preliminary estimates show that considerable additional funding and greater political commitment to maternal, newborn and child health are needed to achieve universal coverage of the full package of interventions in the 68 Countdown countries.
- Official development assistance (ODA):
 - ODA for maternal, newborn and child health increased between 2003 and 2007 but remains far below needed levels. Only 31% of all ODA for health was allocated to maternal, newborn and child health in 2007. Family planning received less funding in 2007 than in 2003.
 - ODA is not always targeted to countries with the greatest need. Achieving

- MDGs 4 and 5 will require donors to improve their funding and allocation practices.
- National resources: although ODA is important, national resources are a much larger share of maternal, newborn and child health funding. Tracking government and nongovernment spending at the country level is essential so that policy-makers can allocate adequate resources for women's and children's health.

Action now

- All countries should:
 - Identify gaps in coverage and quality of care along the continuum of care for maternal, newborn and child health.
 - Improve the delivery of essential interventions and packages.
 - Identify inequities in coverage—by geographic area, ethnic group, income and the like—and initiate actions to provide universal coverage of essential interventions and packages.
 - Increase resource allocations for reproductive, maternal, newborn and

- child health services, ensuring that interventions and programmes are sufficiently funded.
- Other *Countdown* partners should work together with countries to:
 - Advocate for increased funding for reproductive, maternal, newborn and child health through innovative mechanisms and ensure that funding is predictable, consistent and responsive to national needs and plans.
 - Support country efforts to improve data collection and analysis by strengthening health information and vital registration systems as well as by undertaking additional surveys to measure mortality, coverage and funding.
 - Invest in implementation research to identify effective strategies for delivering proven interventions and quantify their impact.
 - Maximize financial and technical support for large-scale implementation of priority strategies and interventions.
 - Encourage the development and use of mechanisms for holding key actors accountable for fulfilling their commitments.





About Countdown to 2015



Countdown to 2015: Tracking Progress in Maternal, Newborn and Child Survival

- is a global movement of academics, governments, international agencies, health care professional associations, donors and nongovernmental organizations, with *The Lancet* as a key partner.
- uses country-specific data to stimulate and support country progress towards achieving the health-related MDGs.
- focuses on coverage of effective interventions for maternal, newborn and child health and coverage determinants, including health systems and policies, financial flows and equity.


Countdown addresses multiple MDGs:

- MDG 4 to reduce child mortality.
- MDG 5 to improve maternal health.
- MDG 1 to eradicate extreme poverty and hunger, specifically by addressing nutrition with a focus on infant and young child feeding.
- MDG 6 to combat HIV/AIDS, malaria and other diseases.
- MDG 7 to ensure environmental sustainability, through tracking improved access to safe water and improved sanitation.

Countdown tracks progress in the 68 countries where more than 95% of all maternal and child deaths occur (map 1).

Countdown focuses on countries (figure 1).

Sample country profile Contains to the country profile Contains



A key element of *Countdown* work is country profiles that bring together information on coverage levels for interventions proven to improve maternal, newborn and child health as

well as relevant demographic, epidemiological, policy, health systems and financing indicators (box 1). The profiles are updated every two to three years with new data and estimates.

BOX 1

Countdown: the best evidence to inform the right decisions

Scope of Countdown work

Countdown focuses on four areas, each addressed by an interdisciplinary technical working group:

- Coverage for interventions proven effective in reducing maternal, newborn and child deaths.
 Coverage is defined as the proportion of the population who can benefit from an intervention who receives it.
- Health systems and policies, which provide an important context for coverage gains.
- Financial flows to maternal, newborn and child health, initially focused on ODA and now including national financing.
- Equity in intervention coverage, focused on the equitable distribution of coverage across socioeconomic quintiles in national populations.

Countdown recognizes the broader set of political, economic, social, technological and environmental determinants of coverage and mortality and incorporates them into analyses where relevant and feasible given available data.

Data sources and methods


The 2008 *Countdown* report describes how the priority *Countdown* countries were identified, the selection of interventions and approaches tracked through *Countdown*, and the coverage indicators associated with each. A full list of *Countdown* indicators and data sources as well as documentation on the methods used to calculate the equity measures, financing gap analysis and ODA estimates is available at http://www.countdown2015mnch.org.

Data quality control

Quality control of the coverage estimates for interventions and approaches effective in reducing maternal, newborn and child mortality is the responsibility of many different groups. *Countdown* supplements these efforts by working closely with the United Nations Children's Fund (UNICEF) and others

responsible for maintaining global databases and conducts additional quality checks to ensure consistency and reliability. Country profiles are shared with ministries of health and UN colleagues prior to publication. Work is under way to address continuing challenges in estimating coverage and associated uncertainty.

Countdown databases within the broader context of maternal, newborn and child survival



New Countdown results expected in 2011

This update reports on data for *Countdown* indicators available as of December 2009. A full *Countdown* report will be released in 2011 and will include:

- Updated estimates for maternal mortality.
- New coverage estimates for many countries drawing on recent national surveys.
- First results from in-depth analyses conducted by all technical working groups.
- Findings from cross-cutting analyses addressing priority questions.
- New ODA estimates for maternal, newborn and child health for 2008, financing for family planning, and domestic spending on maternal, newborn and child health for African and Asian Countdown countries.

Progress towards Millennium Development Goals 4 and 5




Progress towards MDGs 4 and 5 is inextricably linked: improving maternal health will lead directly to reductions in deaths among newborns and young children. There are also shared challenges in improving health services across the continuum from pre-pregnancy through pregnancy, childbirth, the postnatal period, and childhood. This section focuses on trends in mortality; later sections look more in depth at intervention coverage and the related areas of health systems, financial flows and equity.

Good news! There has been huge progress in reducing deaths among children under age 5 worldwide.¹ And yet the opportunity to save children's lives has never been greater. Of the unacceptable burden of nearly 9 million deaths of children under age 5 that occur a year, at least two-thirds of them could be prevented using proven, affordable interventions.² Table 1 shows country-specific progress towards MDG 4, including the estimated under-five mortality rates for 1990, 2000 and 2008, the average annual rate of reduction for 1990–2008 and its trend for 1990–2000 and 2000–2008, and a summary assessment of progress.

Of the 68 *Countdown* countries, 19 are on track to achieve MDG 4 (figure 2), and 17 of those have reduced mortality by at least half. In 47 *Countdown* countries the rate of change in progress over 2000–2008 increased compared with the 1990s. The annual average rate of reduction rose more than 4 percentage points in Azerbaijan, Botswana, China, Ghana, Lesotho, Rwanda and Swaziland. However, further absolute gains are needed to achieve the goal, except in Azerbaijan and Botswana, two countries that are on track.

Much remains to be done. In 12 countries progress has slowed since 2000; some are on track now but may not be if these trends continue (including Guatemala, Indonesia and



the Philippines). National governments and their development partners must stay committed to child survival to prevent reversals in progress and because further gains are harder to achieve as mortality rates fall.

Mortality is not being reduced uniformly. Just over 40% of child deaths now occur in the first month of life. The growing concentration of child deaths in the newborn period is linked to maternal health and survival (box 2). Hundreds of thousands of women die each year because of complications related to pregnancy and childbirth. For every woman who dies, approximately 20 others suffer injuries, infection and disabilities, resulting in millions of women experiencing adverse pregnancy outcomes.

Country progress towards Millennium Development Goal 4

	Under-five mortality rate (per 1,000)			Average annual rate of reduction (%)	Summary	Average annual rate of reduction (%)		
Country or territory	1990	2000	2008	1990-2008	assessment of progress ^a	1990-2000	2000-2008	Direction of change
Afghanistan	260	257	257	0.1	No progress	0.1	0.0	4
Angola	260	239	220	0.9	No progress	0.8	1.0	†
Azerbaijan	98	69	36	5.6	On track	3.5	8.1	↑
Bangladesh	149	91	54	5.6	On track	4.9	6.5	†
Benin	184	144	121	2.3	Insufficient progress	2.5	2.2	+
Bolivia	122	86	54	4.5	On track	3.5	5.8	†
Botswana	50	81	31	2.7	On track	-4.8	12.0	†
Brazil	56	34	22	5.2	On track	5.0	5.4	†
Burkina Faso	201	188	169	1.0	Insufficient progress	0.7	1.3	1
Burundi	189	178	168	0.7	No progress	0.6	0.7	↑
Cambodia	117	106	90	1.5	Insufficient progress	1.0	2.0	1
Cameroon	149	147	131	0.7	No progress	0.1	1.4	↑
Central African Republic	178	181	173	0.2	No progress	-0.2	0.6	†
Chad	201	205	209	-0.2	No progress	-0.2	-0.2	-
China	46	36	21	4.4	On track	2.5	6.7	†
Congo	104	116	127	-1.1	No progress	-1.1	-1.1	_
Congo, Dem. Rep. of the	199	199	199	0.0	No progress	0.0	0.0	_
Côte d'Ivoire	150	138	114	1.5	Insufficient progress	0.8	2.4	↑
Djibouti	123	106	95	1.4	Insufficient progress	1.5	1.4	+
Egypt	90	47	23	7.6	On track	6.5	8.9	†
Equatorial Guinea	198	168	148	1.6	Insufficient progress	1.6	1.6	
Eritrea	150	89	58	5.3	On track	5.2	5.4	†
Ethiopia	210	148	109	3.6	Insufficient progress	3.5	3.8	†
Gabon	92	87	77	1.0	Insufficient progress	0.6	1.5	†
Gambia	153	131	106	2.0	Insufficient progress	1.6	2.6	†
Ghana	118	111	76	2.4	Insufficient progress	0.6	4.7	<u>†</u>
Guatemala	77	47	35	4.4	On track	4.9	3.7	+
Guinea	231	185	146	2.5	Insufficient progress	2.2	3.0	†
Guinea-Bissau	240	218	195	1.2	Insufficient progress	1.0	1.4	↑
Haiti	151	109	72	4.1	On track	3.3	5.2	·
India	116	94	69	2.9	Insufficient progress	2.1	3.9	<u> </u>
Indonesia	86	56	41	4.1	On track	4.3	3.9	J
Iraq	53	48	44	1.0	Insufficient progress	1.0	1.1	†
Kenya	105	128	128	-1.1	No progress	-2.0	0.0	· •
Korea, Dem. Rep.	55	55	55	0.0	No progress	0.0	0.0	
Lao People's Democratic Republic	157	86	61	5.3	On track	6.0	4.3	+
Lesotho	101	109	79	1.4	Insufficient progress	-0.8	4.0	†
Liberia	219	174	145	2.3	Insufficient progress	2.3	2.3	
Madagascar	167	132	106	2.5	Insufficient progress	2.4	2.7	
Malawi	225	162	100	4.5	On track	3.3	6.0	<u>'</u>
Mali	250	217	194	1.4	Insufficient progress	1.4	1.4	
Mauritania	129	122	118	0.5	No progress	0.6	0.4	+
Mexico	45	26	17	5.4	On track	5.5	5.3	1
Morocco	45 88	26 54	36	5.4	On track	4.9	5.3	*
Mozambique	249	183	130				4.3	
•				3.6	Insufficient progress	3.1		1
Myanmar	120 142	107	98	1.1	Insufficient progress On track	1.1	1.1	_
Nepal Niger	305	85	51 167	5.7		5.1	6.4	1
minut	305	227	167	3.3	Insufficient progress	3.0	3.8	1

(continued)

TABLE 1 (continued)

Country progress towards Millennium Development Goal 4

	Unde	Under-five mortality rate (per 1,000)			Summary	Average annual rate of reduction (%)		
Country or territory	1990	1990 2000 2008 1990-2008		1990–2008	assessment of progress ^a	1990-2000	2000–2008	Direction of change
Pakistan	130	108	89	2.1	Insufficient progress	1.9	2.4	1
Papua New Guinea	91	77	69	1.5	Insufficient progress	1.7	1.4	+
Peru	81	41	24	6.8	On track	6.8	6.7	↓
Philippines	61	36	32	3.6	On track	5.3	1.5	+
Rwanda	174	186	112	2.4	Insufficient progress	-0.7	6.3	†
Senegal	149	131	108	1.8	Insufficient progress	1.3	2.4	†
Sierra Leone	278	252	194	2.0	Insufficient progress	1.0	3.3	†
Somalia	200	200	200	0.0	No progress	0.0	0.0	_
South Africa	56	73	67	-1.0	No progress	-2.7	1.1	†
Sudan	124	115	109	0.7	No progress	0.8	0.7	+
Swaziland	84	124	83	0.1	No progress	-3.9	5.0	†
Tajikistan	117	94	64	3.4	Insufficient progress	2.2	4.8	†
Tanzania, U. Rep.	157	139	104	2.3	Insufficient progress	1.2	3.6	†
Togo	150	122	98	2.4	Insufficient progress	2.1	2.7	1
Turkmenistan	99	71	48	4.0	On track	3.3	4.9	†
Uganda	186	158	135	1.8	Insufficient progress	1.6	2.0	↑
Yemen	127	98	69	3.4	Insufficient progress	2.6	4.4	↑
Zambia	172	169	148	0.8	No progress	0.2	1.7	†
Zimbabwe	79	102	96	-1.1	No progress	-2.6	0.8	↑

a. "On track" indicates that the under-five mortality rate for 2008 is less than 40 per 1,000 or that it is 40 or more with an average annual rate of reduction of 4% or higher for 1990–2008; "insufficient progress" indicates that the under-five mortality rate for 2008 is 40 or more with an average annual rate of reduction of 1%–3.9% for 1990–2008; "no progress" indicates that the under-five mortality rate for 2008 is 40 or more with an average annual rate of reduction of less than 1% for 1990–2008.

Source: Mortality rates, Inter-agency Group for Child Mortality Estimation, 2009, www.childmortality.org; average annual rate of reduction and summary of progress, UNICEF 2009a.



BOX 2

New data and continuing challenges in achieving MDG 5 and improving the lives of women

The maternal mortality ratio, the number of maternal deaths per 100,000 live births, is the most common measure of the magnitude of maternal mortality and is a progress indicator for MDG target 5.A, which aims at reducing the ratio by three quarters over 1990–2015, implying an average annual rate of reduction of 5.5%.

New global and regional maternal mortality estimates were set to be released after this *Countdown* update went to press. Updated country estimates will be available in fall 2010 after extensive in-country consultation.

The work on regular updating of the estimates reflects the international community's commitment to continuously improve assessment of the maternal mortality burden but also highlights the urgent need to invest in building country health information systems to monitor maternal mortality. The World Health Organization (WHO), UNICEF, the United Nations Population Fund (UNFPA) and the World Bank work closely with the United Nations Population Division, academic experts and countries to regularly update global, regional and country estimates of maternal mortality. There are preliminary indications of global progress, with some countries having achieved significant declines.¹ A recent academic analysis using alternative statistical assumptions found an annual rate of reduction of 1.3% over 1990–2008,² well short of the 5.5% needed to attain the MDG target.

Measuring maternal mortality remains a challenge. Identifying a maternal death requires accurate data on the deaths of women of reproductive age, including cause of death, pregnancy status and the time of death in relation to pregnancy or childbirth. These data are often missing, misclassified or underreported, particularly in low- and middle-income countries that lack fully functioning vital registration systems and where many women deliver at home. Early pregnancy deaths are especially difficult to identify. The weakness of many developing country health information systems

results in the use of statistical modelling to develop maternal mortality estimates, which better indicate the order of magnitude of the problem. These estimates are subject to considerable uncertainty and vary with the assumptions and methods used. Important limitations of the maternal mortality ratio include:

- The maternal mortality ratio reflects only the risk of death once pregnant and misses the cumulative mortality risk associated with the number of pregnancies a woman has during her reproductive years.
- The maternal mortality ratio is difficult to measure, has large uncertainty bounds and must be interpreted cautiously—hence the need for real-time monitoring and surveillance of maternal deaths.
- The maternal mortality ratio focuses narrowly on mortality and may result in a lack of attention to the millions of women who suffer from "nearmiss" events and short- and long-term pregnancyrelated illnesses.³

Preventing maternal mortality and the millions of pregnancy-related disabilities each year will require concentrated efforts to improve coverage of comprehensive family planning programmes and antenatal, childbirth, emergency obstetric and postnatal care—all indicators tracked in *Countdown*. Improving access to safe abortion care in countries where abortion is legal is also essential for reducing maternal deaths. Coverage of a skilled attendant at birth is a progress indicator for MDG target 5.A and is a sensitive measure of health system strength. Over the past two decades coverage of a skilled attendant at birth has improved in all regions, with considerable gains in North Africa and South-East Asia (see figure 10).4

Notes

- 1. WHO, UNICEF, UNFPA, and World Bank 2007.
- 2. Hogan and others 2010.
- 3. Filippi 2006.
- 4. UNICEF global databases, November 2009.

Causes of maternal and child deaths



What causes the 8.8 million child deaths each year?


New estimates of child deaths for 2008 show that pneumonia, diarrhoea and malaria remain the highest causes worldwide, together accounting for 41% of deaths (figure 3). More than 40% of child deaths occur in the neonatal period, and progress in reducing deaths has been slower for newborn deaths than for deaths among children ages one month to five years. Undernutrition contributes to more than one-third of child deaths.³ The majority of these deaths can and must be prevented by increasing coverage for known, affordable and effective interventions.


The country profiles highlight important regional and country variations in these causes. For example, estimates for Africa indicate that 29% of all child deaths occur in the neonatal period and that 49% of deaths after this period are due to pneumonia, diarrhoea or malaria.⁴ In contrast, estimates for South East Asia indicate that about

54% of child deaths occur in the neonatal period and that about 26% of postneonatal deaths are due to pneumonia, diarrhoea or malaria.

Global distribution of maternal causes of death


A maternal death is defined as the death of a woman while pregnant or within 42 days of termination of pregnancy, regardless of the site or duration of pregnancy, from any cause related to or aggravated by the pregnancy or its management. New estimates show that the leading causes of maternal deaths are haemorrhage and hypertension, which together account for more than half of maternal deaths (figure 4). Indirect causes, which include deaths due to conditions such as malaria, HIV/AIDS and cardiac diseases, account for about onefifth of maternal deaths. Regional estimates show that haemorrhage and hypertension are among the top three causes of deaths in both South Asia and Sub-Saharan Africa, where the





majority of maternal deaths occur. This is in contrast to developed countries, where other direct causes—for example, those related to complications of anaesthesia and caesarean sections—are the leading cause of death, reflecting global disparities in access to needed obstetrical care.

The categories of maternal deaths are based on a new classification system developed by WHO that considers obstructed labour and anaemia to be contributing conditions rather than direct causes. Deaths related to these two conditions are now classified within the categories of haemorrhage or sepsis.



Social determinants of maternal, newborn and child health



In the words of the WHO Commission for Social Determinants of Health (2008), social determinants of maternal, newborn and child health "... are the conditions in which people are born, grow, live, work and age, including the health system. These circumstances are shaped by the distribution of money, power and resources at global, national and local levels, which are themselves influenced by policy choices. The social determinants of health are mostly responsible for health inequities—the unfair and avoidable differences in health status seen within and between countries."

Poverty and inequity are underlying contributors to many maternal, newborn and child deaths, and evidence shows that poor households have more than twice the risk of mortality of wealthy households. Poverty affects maternal, newborn and child health through a range of mechanisms. Poor diets and food insecurity increase the risk of illness and undernutrition (box 3); poor environmental conditions contribute to inadequate housing, water and sanitation; and family poverty reduces care-seeking and access to information and health care services. Poverty and lack of access to care can be compounded by conflict,

вох з

Undernutrition: a risk for women and children

Undernutrition affects mortality and ill-health along the entire continuum of care from pre-pregnancy to early childhood.

Undernutrition, the result of poor dietary quality and inadequate intake of micronutrients as well as low energy intake, contributes to at least one-third of child deaths. Stunting, or low height for age, is a particularly important *Countdown* indicator because it reflects longer term nutritional deficiencies with implications for growth and development of children now and in future generations. Child undernutrition and infectious diseases are synergistic and cyclical, posing a major threat to child survival.

Maternal short stature and iron deficiency anaemia, which can increase the risk of death of the mother at delivery, contribute to at least 20% of maternal deaths. Maternal undernutrition also increases the probability of low birth weight, which in turn increases the probability of neonatal deaths due to infections and asphyxia. Measures of maternal undernutrition will be tracked by *Countdown* beginning in the 2011 report.

Two-thirds of the world's children affected by stunting live in just 10 *Countdown* countries

Country	Stunting prevalence (%)	Number Share of developing children (thousands) (%)
India	48	60,788 31.2
China	22	12,685 6.5
Nigeria	41	10,158 5.2
Pakistan	42	9,868 5.1
Indonesia	37	7,688 3.9
Bangladesh	43	7,219 3.7
Ethiopia	51	6,768 3.5
Congo, Dem. Rep. of the	46	5,382 2.8
Philippines	34	6,317 1.9
Tanzania, U. Rep.	44	3,359 1.7
Total		65.5

Source: UNICEF 2009b.

Source: Stewart, Dewey, and Ashorn 2010; Victora and others 2008; Black and others 2008; UNICEF 2009b.

population displacement and emergencies such as floods and drought. Recent analyses indicate that maternal, newborn and child health can be negatively affected by high burdens of noncommunicable diseases that increase the likelihood of catastrophic expenditures at the household level. Maternal, newborn and child health is also influenced by gender discrimination, low levels of female education, few incomeearning opportunities for women and other societal factors affecting women's empowerment. Further, the death of a mother increases the risk that her children will die.

A range of measures are available to address social determinants of health. Expanding educational programmes, introducing gender-based affirmative action policies and other programmes to achieve MDG3, implementing laws supportive

of human rights and improving living and working conditions (for example, improving access to clean water and adequate sanitation) can all make a difference. Good governance and oversight of health systems can positively influence maternal, newborn and child health in difficult circumstances. In addition, tackling the inequitable distribution of power, money and resources should be a priority. Other successful approaches include addressing financial barriers to care by, for example, reducing or eliminating user fees or introducing targeted conditional cash transfer schemes. Women's support groups have been shown to contribute measurable improvements in maternal, newborn and child health as well as mental health, suggesting that such strategies can be employed synergistically with health sector reforms to improve women's empowerment and decision-making.



Coverage along the continuum of care



Countdown tracks coverage along a continuum of care from pre-pregnancy and childbirth through childhood up to age 5, highlighting missed opportunities for the delivery of lifesaving interventions. Median coverage levels for 20 Countdown interventions are summarized in figure 5 but they mask important variations in levels and progress at the country level.

Coverage gaps


The 2010 *Countdown* results show important gaps in coverage for three groups of interventions:

 Interventions immediately surrounding birth, such as the presence of a skilled attendant, ensuring the early initiation of breastfeeding and

- an early postnatal visit to check on the health of the mother and newborn.
- Interventions that require 24-hour access to a skilled health provider, such as treatment of childhood pneumonia, diarrhoea and malaria.
- Interventions introduced only recently, such as intermittent preventive treatment for malaria during pregnancy, or recently scaled-up interventions, such as the use of insecticide-treated nets.

Equity gaps

Data on the disparities in coverage between the poorest and the least-poor, shown later, highlight the poor-rich gap in access to these essential



a. Target coverage value is not 100%.

Source: Prevention of mother-to-child transmission of HIV/AIDS, UNICEF, Joint United Nations Programme on HIV/AIDS (UNAIDS) and WHO; immunization rates, WHO and UNICEF; postnatal visit for mother, Saving Newborn Lives analysis of Demographic and Health Surveys; improved water and sanitation, WHO and UNICEF Joint Monitoring Programme 2010; all other indicators, UNICEF Global Databases, November 2009, based on Demographic and Health Surveys, Multiple Indicator Cluster Surveys and other national surveys.

lifesaving interventions, including family planning services.

Quality gaps

Coverage estimates for service delivery contacts—such as antenatal care, skilled attendant at birth and postnatal visits for the mother—do not address the quality of that contact or whether it provided needed interventions such as active management of the third stage of labour or counselling on family planning. Quality assessments of such

services are an essential part of sound programme management.

Data gaps

Countdown highlights data gaps that must be addressed to improve the ability of countries to make informed decisions on how to accelerate progress towards MDGs 4 and 5. For example, only 23 Countdown countries have data available on postnatal care for women, and six have data on postnatal care for newborns.





Every pregnancy wanted

Addressing adolescent reproductive health an essential part of the continuum of care


Newly included in this *Countdown* update are estimates of the adolescent birth rate, defined as the annual number of births to women ages 15–19 per 1,000 women in that age group. This is a progress indicator for MDG target 5.B for achieving universal access to reproductive health. Adolescent fertility is high in many *Countdown* countries (figure 6), which means that many young women face an elevated risk of maternal death and disability. Newborns and infants of adolescent mothers are also at higher risk of low birth weight and mortality.

Analysis of 23 *Countdown* countries in Sub-Saharan Africa with two consecutive Demographic and Health Surveys since 2000 shows at least a 10% drop in the adolescent fertility rate in 18 of them. In the majority of these countries, the declines are primarily among women from wealthier households, those living in urban areas and those with higher education levels.

Increasing access to family planning


Reducing unwanted pregnancies reduces overall births, including those among adolescent women, and therefore reduces maternal deaths and unsafe abortions. The impact of birth spacing on newborn and child survival is also important. Countdown tracks both the contraceptive prevalence rate—the percentage of women married or in union ages 15–49 who are practising, or whose sexual partners are practising, any form of contraception—and the unmet need for contraception—the percentage of married women who do not want a child or who want to postpone their next pregnancy but are not using any contraception (figure 7). Both are progress indicators for MDG target 5.B.

Trends in family planning coverage have been highly variable across countries. The small increase in the median coverage for the 42 countries with data on contraceptive prevalence from around 2000



Disparities in family planning coverage and need are wide

Unmet need for family planning, *Countdown* countries with the highest and lowest contraceptive prevalence rates, various years since 2000 (%)



a. Refers to unweighted median of all *Countdown* countries with data available (49 countries for family planning and 66 countries for contraceptive prevalence rate).


Source: Demographic and Health Surveys, Multiple Indicator Cluster Surveys and other national household surveys.

and around 2008 mask important increases and lack of progress in individual countries (figure 8). For example, Madagascar (box 4), Rwanda and Swaziland all increased coverage by more than 20 percentage points, but 12 other countries showed no change or a decrease in coverage, with Central African Republic and Togo seeing declines of

FIGURE 8

Progress in coverage for family planning since 2000 has been mixed

Coverage for family planning, contraceptive prevalence rate and trends for *Countdown* countries with the highest and lowest contraceptive prevalence rates; 2008 (%)



a. Refers to unweighted median of 42 *Countdown* countries with data available for both time periods.

Source: Demographic and Health Surveys, Multiple Indicator Cluster Surveys and other national household surveys.

9 percentage points and Democratic Republic of Congo a decline of 11 percentage points. Decreased ODA to family planning from 1990 to 2007, among other factors, likely contributed to this lack of progress (see figure 22). There are also disparities in family planning coverage within countries, with lower coverage among women in poorer households and among adolescents relative to older women.



BOX 4


What can we learn from Madagascar's family planning success?

Contraceptive prevalence in Madagascar rose from 5.1% in 1992 to 29% in 2008–09, according to Demographic and Health Surveys (see figures). The Madagascar Family Planning Program attributes this success to three aspects of the programme:


- Leadership and policy
 - Strong leadership at the highest level (such as former President Marc Ravalomanana).
 - A target included at the highest level in the national development plan (Madagascar Action Plan target of 30% contraceptive prevalence in 2012).
 - Policy of providing contraceptives free in all public health facilities.
- Community involvement
 - Effective multisectoral collaboration with involvement of local authorities (17,433).
 Fokontany chiefs initiated and sensitized on family planning through a film and brochures).
 - National campaign of integrated activities for family planning, adolescent reproductive health and HIV/AIDS in 81% of public schools.
 - 3,000 women leaders initiated and sensitized on family planning.
 - Community health workers trained to provide family planning via injectable contraceptives.
- Programme management
 - Integration of family planning programme in all functional public health facilities, including 50 voluntary counselling and testing centres (all public and functional).
 - Family planning commodity security assured through coordination by a logistic committee, monitoring and periodic surveys.

The prevalence of modern contraception among married women in Madagascar has risen consistently since 1992

Contraceptive prevalance rate (%)



The number of women using family planning has accelerated since 1992



Source: Demographic and Health Surveys.



Every birth safe

All women and their newborns need skilled care at birth and access to emergency care when complications develop. Outcomes around birth are a sensitive marker of the strength of health systems, including the quality of available care (figure 9). Life-threatening complications at birth require rapid response. Postpartum haemorrhage can kill a mother in a few hours, and a newborn who is not breathing at birth will be dead within minutes.

Addressing current global gaps for care at birth is critical to achieving MDGs 4 and 5. When mothers die during childbirth, it is rare for the newborn to survive, and the risk of mortality increases for any young children left behind in the household.


Coverage gap for care at birth

Some *Countdown* countries have made good progress in increasing the proportion of women attended by a skilled provider during childbirth, a progress indicator for MDG target 5.A. Three countries—Burkina Faso, Pakistan and Rwanda—had gains of more than 20 percentage points from around 2000 to around 2008, and 10 had gains of more than 10 percentage points since 1990 (figure 10). Gains were uneven, however, with 11 countries showing no progress. More effort is needed to ensure all pregnant women and newborns have access to a skilled provider.

Coverage gap for emergency obstetric care

Life-threatening complications during labour and delivery are often unpredictable and unpreventable. All pregnant women must have access to skilled care at birth (box 5) and a guarantee that basic or emergency obstetric care services are accessible when needed. Availability of such services is low in many *Countdown* countries (see page 29). Caesarean section coverage rates below 5% signal a lack of access to emergency obstetric care and indicate human resources and other health systems challenges⁷; 33 of the 51 *Countdown* countries with data since 2000 had rural rates below 5%, and 4—Burkina Faso, Chad, Ethiopia and Niger—had rural rates below 1%.

The period surrounding birth accounts for a high proportion of deaths



Source: Adapted from Lawn and others (2009) using data on stillbirths from Stanton and others (2006), data on intrapartum stillbirths from Lawn, Shibuya, and Stein (2005), data on neonatal deaths from Black and others (forthcoming) and data on maternal time of death from Li and others (1996).


Quality gaps in care before, during and immediately after childbirth

Antenatal, delivery and postnatal care are service contact points and offer opportunities for the provision of effective interventions that can prevent illness and save lives. Median coverage for four or more antenatal care visits was 50% across the 51 *Countdown* countries with data since 2000, ranging from 6% in Somalia to 89% in Brazil. Contact with a trained service provider during pregnancy provides the opportunity for delivery of numerous proven interventions that improve outcomes for mother and newborn, including treatment of hypertension to prevent eclampsia, tetanus

Progress in increasing the proportion of women attended by a skilled provider at

birth has varied

Coverage of live births attended by a skilled health worker, Countdown



Source: UNICEF Global Databases, November 2009, based on Demographic and Health Surveys, Multiple Indicator Cluster Surveys and other national household surveys.

immunization, intermittent preventive treatment for malaria and distribution of insecticide-treated nets, prevention of mother-to-child transmission of HIV, micronutrient supplementation, and counselling on family planning and birth preparedness. UNICEF, UNFPA and WHO recommend at least four antenatal care visits at key stages through pregnancy. The number of visits alone will not save lives, of course, unless high-quality, effective interventions are delivered during visits. Work is continuing on the

BOX !

Strategies for increasing coverage for skilled birth attendant at birth

Improving coverage of skilled attendant at birth requires strategies that address supply- and demandside barriers to care and service quality. Some countries, such as Indonesia, are addressing supplyside barriers by training a new cadre of community midwives and bringing care into the home and local community. Other countries, such as Mozambique, are using task-shifting to enable midwives and other nonphysician clinicians to provide essential interventions, including caesarean sections. Subnational examples of progress in India include public-private partnerships to increase the number of private obstetricians delivering services to the rural poor in Gujarat, conditional cash transfers and a remuneration mechanism for community health workers.¹ These initiatives need to be assessed for impact and for the feasibility of being scaled up. On the demand side, reducing financial barriers is a common feature of success. Ghana recorded a rapid increase in facility births linked to the introduction of a national insurance scheme and new policies guaranteeing free care at birth.² Rwanda's "Paying for Performance" strategy increased institutional deliveries by providing financial incentives to providers to increase the use and quality of care.3

Notes

- 1. Mavalankar and others 2009; Lee and others 2009; India 2005; Devadasan 2008
- 2. Witter, Armar-Klemsesu, and Graham 2009.
- 3. Basinga 2010.

measurement challenges associated with assessing coverage for individual interventions and service quality during antenatal care visits.

Not all women who have contact with a health provider during childbirth and in the immediate postnatal period receive the range of interventions that are needed (such as active management of the third stage of labour with the delivery of oxytocin to prevent post-partum haemorrhage). This quality gap is a missed opportunity to improve maternal and newborn health and reduce stillbirths. Postnatal care for mothers and newborns is another gap: data are lacking for many countries (45 of the 68 have no data), coverage is low in the 23 countries with data available (median coverage of 38%), and effective interventions are often not provided. Only six *Countdown* countries have data on postnatal care for the newborn, and the median coverage is low (4%).



Every newborn and child healthy


Solutions and innovations to save newborn lives

Risk of death is high for both mother and newborn in the first few days of life. There has been an explosion of interest and research in preventing newborn deaths, but some newer interventions are not yet reflected in *Countdown* tracking. Two examples are antenatal steroids—a high-impact, evidence-based intervention delivered during preterm labour that has been associated with a 53% reduction of newborn deaths due to preterm birth complications —and kangaroo mother care—a simple technique where the newborn is kept close to the mother's body in front, providing warmth, increased feeding, reduced infections and more rapid recognition of illness. New evidence shows that hospital-based kangaroo mother care

can reduce deaths for newborns under 2,000 grams (almost all preterm) by 51%.¹⁰

Evidence on the importance of providing postnatal care within two days of delivery has led to a joint WHO–UNICEF statement calling for broader implementation and scaling up.¹¹ Simple interventions such as early initiation and exclusive breastfeeding (figure 11), keeping the newborn warm, hygienic cord and skin care have the potential to reduce a large number of newborn deaths.

Other known, proven interventions to protect the lives of newborns and young infants are tracked by *Countdown* to determine progress in coverage at the country level. Rates in 2010 show some important gains, but many missed opportunities.



Source: UNICEF Global Databases, November 2009, based on Demographic and Health Surveys, Multiple Indicator Cluster Surveys and other national surveys.

Progress in preventing major childhood infectious diseases

Countdown results highlight progress on a country-by-country basis in preventing the infectious diseases responsible for the majority of child deaths. There has been important progress in combating malaria through increased use of insecticide-treated nets in malaria-endemic countries (figure 12) and in preventing mother-to-child transmission of HIV (box 6). More work needs to be done, however, for countries to reach the Roll Back Malaria target of 80% coverage of insecticide-treated nets by 2010 and universal coverage of prevention of mother-to-child transmission for HIV positive pregnant women.

Greater attention to improved water and sanitation can prevent diarrhoea

MDG target 7.C on environmental sustainability is to halve, by 2015, the proportion of people without

sustainable access to safe drinking water and basic sanitation. Median coverage in the 65 *Countdown* countries with data available since 2000 is 71% for use of an improved drinking water source and 41% for improved sanitation (figure 13). More efforts are needed to promote better hygiene and to ensure that adequate drinking water and sanitation are available to all in order to reduce child deaths from diarrhoea.

Improving infant and young child feeding practices will save lives

Available evidence demonstrates that child growth and development are optimized when:

- Breastfeeding is initiated within one hour of birth.
- Exclusive breastfeeding is continued up to age six months
- Complementary feeding with safe and age-appropriate solid, semi-solid or soft foods is started at age six months.

BOX 6

Preventing mother-to-child transmission of HIV

HIV accounts for a relatively small proportion of deaths among children under age 5 across the *Countdown* countries as a whole. But in a subset of 15 high-HIV-burden *Countdown* countries (those with prevalence of 5% or higher), it continues to be a major threat to survival and child development (see table). Preventing HIV infection in women and children requires a strategy across the continuum of care, integrating:

- Interventions directed at reducing infection among young people with access to information and testing.
- Interventions to meet the family planning needs of women living with HIV.
- Antiretroviral therapy where needed.
- Safe practices during childbirth.
- Guidance for selecting safe and optimal infantfeeding options to prevent mother-to-child transmission of HIV.
- Provision of antiretroviral regimens for the prevention of mother-to-child transmission of HIV.
- Scaling up early infant diagnosis to ensure prompt and effective treatment of infections.

There have been dramatic increases in prevention of mother-to-child transmission coverage in 9 of the 15 high-HIV-burden *Countdown* countries. Cameroon and the Central African Republic saw more modest gains, and Botswana (already at 95% coverage) and Congo saw limited gains (Malawi and the United Republic of Tanzania do not have data for 2008). These results


demonstrate what is possible when both commitment and resources are focused on reaching a target population with a specific intervention.

Share of HIV-infected women ages 15–49 who received antiretroviral regimens for prevention of mother-to-child transmission, high-HIV-burden *Countdown* countries, 2006 and 2008

	2006		2008		
Country	Point estimate	Range estimate	Point estimate	Range estimate	
Botswana	95	95->95	>95	75->95	
Cameroon	22	18-30	28	20-53	
Central African Republic	18	16-20	23	16-44	
Gabon	4	3-5	35	22-70	
Kenya	48	42-59	56	37->95	
Lesotho	17	15-18	57	43->94	
Malawi	14	12-16	_	41->95	
Mozambique	13	11–15	42	26-93	
South Africa	50	43-60	73	53->95	
Swaziland	62	57-69	>95	87->95	
Tanzania, U. Rep.	15	14-16	_	53->95	
Uganda	25	22-28	50	36-95	
Zambia	35	31-39	59	43->95	
Zimbabwe	17	16-19	36	26-64	

— is not available.

Source: 2006 data, WHO and UNICEF (2007), with the denominators derived from unpublished UNAIDS-WHO estimates; 2008 data, UNICEF, UNAIDS, WHO, and UNFPA 2009.





Note: For each country, the left bar shows the most recent year with data on coverage values and the right bar shows data for a previous year. Source: UNICEF Global Databases, November 2009, based on Demographic and Health Surveys, Multiple Indicator Cluster Surveys and other national surveys.

Some *Countdown* countries are progressing in these areas, and 12 have increased exclusive breastfeeding rates by 20 percentage points or more (see figure 11). But most *Countdown* countries have much room for improvement (figure 14). Current median coverage based on latest available estimates since 2000 are 48% (ranging from 20% to 78%) for early initiation of breastfeeding, 34% (ranging from 1% to 88%) for exclusive breastfeeding among infants less than age 6 months and 66% (ranging from 15% to 93%) for timely introduction of complementary feeding.

Vaccination and vitamin A

Vaccination coverage rates have generally remained high, with median 2008 coverage rates in Countdown countries of 79% for measles vaccination (ranging from 23% in Chad to 99% in Brazil and Turkmenistan) and 83% for diphtheria and tetanus with pertussis (DPT3) vaccination (ranging from 20% in Chad to 99% in Morocco and Peru). Vaccination rates for neonatal tetanus also remain high, with a median of 83% of newborn children considered protected at birth in 2008 (ranging from 47% in Lao PDR to 97% in Sierra Leone). New vaccines for pneumococcal pneumonia and diarrhoea due to rotavirus can build on these delivery successes as they are scaled up in the next few years. Vitamin A supplementation (two doses) estimates for 2008






show a median of 86%, ranging from no vitamin A coverage in Chad and Gabon to 100% coverage in Burkina Faso and Somalia. Chad and Gabon faced major challenges to vitamin A delivery in 2008: Chad experienced a looting of supplies, and Gabon lacked the funding needed to carry out Child Health Days. These two examples are important reminders of the challenges many *Countdown* countries face in sustaining basic services.

and other national surveys.

Coverage of correct treatment for childhood illness remains too low

Progress in care-seeking and case management of common childhood diseases has been very slow. Scaling up case management requires families and communities to be aware of danger signs and to bring children for care. It also requires quality care to be available, which requires adequate human resources and commodities. Unless care is provided close to home, reducing mortality rates will be difficult (box 7).

Correct treatment of diarrhoea includes reducing susceptibility to severe diarrhoea and dehydration through improved nutrition and prompt treatment of watery diarrhoea with oral rehydration salts solution and zinc while continuing to feed the child. Median coverage of correct treatment of diarrhoea in *Countdown* countries was only 42% (figure 15)—a figure that masks variability across countries and



in some instances within countries. There has been rapid policy uptake of "new" low-osmolarity oral rehydration salts and zinc, with 46 *Countdown* countries reporting having adopted such a policy.

Correct treatment of childhood pneumonia and neonatal infections (sepsis and pneumonia) includes antibiotics. This requires a caregiver to recognize signs of illness and seek care from a trained provider. Median coverage of careseeking was only 48% for the 64 *Countdown* countries with data available, while the median coverage of children with suspected signs of pneumonia who actually received an antibiotic was 27% in 35 countries with data.

Correct treatment of childhood malaria requires administration of an effective antimalarial within 24 hours of onset of symptoms.¹³ The current "gold standard" treatment in most malaria-endemic countries is artemisinin-based combination therapies, for which funding and procurement have rapidly increased. Beginning in 2010, *Countdown* will track coverage by type of antimalarial treatment because treatment with chloroquine and other antimalarials is no longer effective in most malaria-endemic countries. Figure 16 shows that tracking coverage by type of antimalarial is important for determining whether children are receiving effective treatment.

Bringing care for sick children closer to home

What evidence is there of the effect of community case management?

Community case management requires trained community health workers to deliver high-impact, curative interventions to children whose families lack access to facility-based care. Recent WHO-UNICEF joint statements summarize the evidence that community health workers can recognize and manage common life-threatening childhood illnesses. 1 The statements cover diarrhoea, pneumonia, malaria and uncomplicated severe acute malnutrition. Several studies report positive outcomes of community case management on pneumonia, including a recent review suggesting a 70% reduction in pneumonia deaths among children under age 5² and others showing the effectiveness of community health worker administration of oral antibiotics for neonatal pneumonia in the absence of referral. Community case management has also been used effectively for malaria (including with artemisinin-based combination therapies) and diarrhoea treatment.

Where is community case management working?

Since the 2008 Countdown report, 11 countries have changed policy to allow community-based management of pneumonia, increasing the total number of Countdown countries in support of community case management to 29. Nepal and Senegal have already scaled up community programmes for management of pneumonia with positive results. Ethiopia and Uganda recently adopted supportive policies and are ready to introduce and rapidly scale up integrated community case management. And India and Malawi now implement integrated management of childhood illness at the community level. Preliminary results from Malawi have shown that health surveillance assistants (governmentpaid, multipurpose extension health workers) can perform an integrated assessment and treat children

appropriately. Families appreciated the proximity and quality of care, and service utilization increased.


What is next for community case management?

Few studies or programmes have systematically evaluated the process and effect of integrated community case management for a comprehensive range of neonatal and childhood illnesses. However, evaluations of the effect of community case management for multiple childhood illness conditions are under way. More work is also necessary to assess a recommended package of services and tasks that a community health worker can deliver reasonably well. For example, would a community health worker providing community case management also be able to provide home-based newborn care? Studies to develop simplified antibiotic regimens for the treatment of neonatal sepsis have commenced and will inform the future role of community health workers in the treatment of severe newborn illness. Two studies from South Asia, for example, present evidence that community health workers can correctly provide treatment for neonatal sepsis with injection gentamicin, but this intervention has not been widely implemented.3 Community case management may be particularly effective in settings where populations are experiencing conflict or natural disasters, but data are lacking.

Countdown countries are tracking the evidence on community case management for newborns carefully, because standard inpatient treatment for seven days for newborn illnesses is not feasible for some families in many of these settings. Including the treatment of uncomplicated severe acute malnutrition in integrated community case management is a possibility.

Notes

- 1. WHO and UNICEF 2004b.
- 2. Theodoratou and others 2010.
- 3. Bang and others 1999; Baqui and others 2009.



The building block framework: health systems and policies to save the lives of women, newborns and children



Coverage of effective interventions is to a large extent the result of the quality and effectiveness by which health services reach people and people's to access them. A well functioning health system comprises several building blocks that have multiple relationships and interactions, with people at the centre. To understand the context in which countries are making efforts to scale up maternal, newborn and child health interventions, the *Countdown* examined progress of key indicators related to each of the building blocks, complementing the information on intervention coverage (a direct function of service delivery).

Health systems are more than building blocks




Source: Alliance for Health Policy and System Research 2010.

Human resources

The challenge: shortages and maldistribution

Only 15 *Countdown* countries (22%) meet the critical threshold of 23 doctors, nurses and midwives per 10,000 people generally considered necessary to deliver essential health services.¹⁴ The shortage is compounded

Median density of doctors, nurses and midwives (per 10,000 people)



Note: Median refers to 26 Countdown countries with data available. Source: WHO 2009.

by uneven geographic distribution within countries.¹⁵

Increased investment in education of health workers, strategies for motivating health workers to remain in underserved areas and effective regulatory frameworks (including those for skills substitution) are among the effective policy options for addressing critical workforce shortages and maldistribution.

One potential solution: task-sharing to supplement services

Ethiopia, Ghana and Pakistan are among the countries addressing workforce shortages and maldistribution challenges through comprehensive strategies, including deployment of health service providers at the community level. The United Republic of Tanzania and Zambia have authorized nonphysician clinicians to carry out certain specialized tasks. More than 90% of caesarean sections in rural areas in Malawi and Mozambique are performed by surgical technicians, with low morbidity and mortality. Malawi and mortality.

Health financing

The challenge: excessive out-of-pocket payments


Median per capita health expenditure in the *Countdown* countries is \$80 (in 2007 international dollars), and only five countries devote at least 15% of their national budgets to health.¹⁹ Only five countries have out-of-pocket expenditure as a percentage of total health expenditure of less than 15%; above this value households may be more vulnerable to catastrophic payments.²⁰

Maternal, newborn and child health services should be available, of good quality and free at the point of delivery in order to remove financial barriers to access and utilization.²¹

Some solutions to move to universal access

A key step towards universal coverage is to move away from out-of-pocket payments through prepayment and risk-pooling schemes.²² Several countries—Mexico, with *Seguro Popular*,²³ and China,²⁴ with the New Rural Cooperative Medical

Scheme—are moving in this direction. In Mali and Rwanda social health insurance schemes are achieving high coverage²⁵ and showing a positive effect on access to priority health services, including maternal, newborn and child health.²⁶ Uganda increased coverage of essential health services, particularly among the poor, by removing user fees.²⁷



Medicines and equipment


The challenge: continuous supply of commodities

Access to medical products, technologies and essential drugs remains erratic in many countries, contributing to low coverage of family planning and poor availability of emergency obstetric care services.

Updated policies on care for maternal, newborn and child conditions need to be accompanied by investment in infrastructure, medicines and supplies, in order to reach universal coverage and make an impact on women's and children's lives.

Possible solutions

Local production of zinc blister packs in Bangladesh combined with health education and promotion through private sector has rapidly increased coverage of zinc use rate in children with diarrhoea to 20%.²⁸ Globally, UNICEF increased procurement of zinc from 20.5 million tablets in 2006 to 158 million tablets in 2008.



Median percentage of emergency obstetric care services considered fully functional, based on assessments of infrastructure, equipment, medicines and skills, 30 *Countdown* countries

Source: WHO, UNFPA, UNICEF, and AMDD 2009.

Health information


The challenge: timely, high-quality information

Decision-makers in most *Countdown* countries do not have the information they need to inform decisions and guide action.²⁹ Vital statistics, including birth registration and maternal death notification, and programme funding are examples of mechanisms that can address barriers to the quality of and access to health services.³⁰

One potential solution: maternal death audit and remedial action

The comprehensive monitoring and evaluation approach adopted in Tamil Nadu, India, which includes stronger information generation and use for decision-making accompanied by maternal death audits and surveillance, has improved maternal and child indicators.³¹

South Africa has also made progress in institutionalizing maternal death audits, which can reduce maternal and perinatal mortality.³²



Source: WHO 2009.

Governance and leadership


The challenge: universal adoption of policies that support maternal, newborn and child health

Too few countries have adopted evidence-based policies to increase access to and quality of essential maternal, newborn and child health services as recommended by WHO–UNICEF joint statements.³³ Just over one-third of *Countdown* countries allow midwives to perform lifesaving interventions, while 46 countries have updated guidelines on management of diarrhoea. Nevertheless, 29 countries have adopted national policies allowing community health

workers to manage pneumonia, compared with 18 in the 2008 *Countdown*. The presence of a national, costed MNCH implementation plan, necessary to estimate and allocate resources efficiently was reported by 41 countries.

Possible solutions

Bangladesh and Nepal have shown that policies that increase access to treatment of diarrhoea and pneumonia in the community are effective in increasing coverage of child health interventions.³⁴ Allowing midwives to perform lifesaving interventions increases access to basic emergency obstetric care services and can reduce maternal mortality.³⁵



Equal care for every mother and child



National data on coverage levels often hide important disparities among population subgroups. The *Countdown* equity analyses include systematic breakdowns of key coverage indicators by wealth quintiles, maternal education, sex of the child, urban/rural residence and region of the country.³⁶

One way to summarize equity analyses is through a mean coverage index. Each country profile includes a figure showing the mean coverage index consisting of an unweighted average of four intervention areas across the continuum of care. Each area includes selected indicators for eight reproductive, maternal, newborn and child interventions: family planning (need for family planning satisfied), maternal and newborn health (at least one antenatal visit and skilled attendant at delivery), immunizations (measles, BCG and DPT3) and curative child care (diarrhoea and pneumonia management: oral rehydration and continued feeding and care seeking for pneumonia).


Figure 18 uses Benin as an example. The mean coverage index of the eight interventions is 73% among children in the richest wealth quintile compared with 41% in the poorest wealth quintile. The top part of figure 18 shows the coverage gap—or how much of an increase is needed to achieve universal coverage with all eight interventions. The gap equals 100% minus the mean coverage index and is much greater for the poor than for the rich.

In all 38 Countdown countries with Demographic and Health Survey data, intervention coverage is substantially higher among mothers and children from better-off households than among those from poor households (figure 17). Countries with similar levels of overall coverage may differ substantially in terms of equity. For example, both Guatemala and Zambia have an overall coverage index of 59%, but in Guatemala mothers and children from households in the poorest quintile show 38% coverage while in Zambia they show 55% coverage. Countries with smaller gaps between rich and poor—such as Bangladesh, Brazil, Egypt, Swaziland and Zambia-should be better studied to understand how they managed to reduce inequalities.

The equity analysis gaps are markedly larger for maternal and newborn interventions than interventions delivered to older children (figure 19). Interventions that are most frequently delivered in fixed health facilities (for example, antenatal, delivery or postnatal care) tend to show greater disparities than those delivered at the community level (for example, vaccinations, vitamin A supplementation or insecticide-treated nets). Family planning interventions, which may be delivered in fixed facilities or at the community level, fall between these two groups in terms of inequalities. Early initiation of breastfeeding shows remarkably small disparities—possibly because it is largely dependent on longstanding cultural practices not yet affected by promotion efforts.

FIGURE 17

The gap in intervention coverage between rich and poor households varies by country



Note: Mean coverage index is based on coverage rates of eight maternal, newborn and child health interventions: met need for family planning, at least one antenatal care visit, skilled attendant at birth, measles vaccination, DPT3 vaccination, BCG vaccination, oral rehydration and continued feeding, and careseeking for pneumonia.

Source: Demographic and Health Surveys.


FIGURE 18 The coverage gap in Benin for eight maternal, newborn and child health interventions decreases as wealth increases Coverage (%) ■ Coverage gap ■ Mean coverage 100 75 50 25 0 Poorest 2nd Richest Wealth quintile

FIGURE 19

Source: Demographic and Health Surveys.

The degree of inequality is markedly larger for maternal and newborn interventions than for those delivered to older children

Average coverage levels of selected reproductive, maternal, newborn and child interventions, poorest and richest wealth quintile, 38 *Countdown* countries with data



Note: Postnatal care refers to postnatal care for all newborn infants. Source: Demographic and Health Surveys.



Country spotlight: Brazil's success in narrowing the gap




Brazil is one *Countdown* country on target to reach the MDGs related to child health and nutrition. Under-five mortality has been falling an average of 5.2% a year since 1990, considerably faster than the 4.4% needed to reach the MDG target. Currently, 22 of every 1,000 children now die before their fifth birthday.³⁷ Underweight prevalence among children under age 5 dropped from 5.7% in 1990 to 2.2% in 2006–2007,³⁸ while stunting fell from 19.9% to 7.1%.


Overall progress has been accompanied by a sharp decline in inequalities (see figures). In 1996 just over 70% of all births to mothers in the poorest socioeconomic quintile received skilled care during childbirth, but by 2007 coverage was universal. Likewise, stunting prevalence among children in the poorest quintile fell from 40% in 1989 to 10% in 2007, while remaining stable at 3%–5% in the richest quintile. These are only two examples among many for which equity improved in the last two decades.

Brazil's success in reducing inequities cannot be attributed to a single factor. Although economic growth has been moderate since 1990, income distribution showed marked improvements in recent years. A nationwide, tax-based unified health system without any user fees was launched in 1988, and geographic targeting has guided the deployment of family health teams of doctors, nurses and community health workers in the poorest areas of the country. As a result, primary health care coverage is now virtually universal, as seen with skilled delivery. In addition, Brazil's conditional cash transfer programmes cover about one third of the population, and multiple integrated health sector initiatives—including immunization, HIV control and breastfeeding promotion activities—have been highly successful.

Perhaps more than any single policy or initiative, the reduction of regional and socioeconomic disparities in health and development has been a central element in Brazil's political agenda for the last 20 years, and it is now starting to bear fruit.



Source: Barros and others 2010.




Source: Monteiro and others 2010.

Country spotlight: Narrowing gender differentials in Bangladesh



Historically, Bangladeshi boys have been more likely to receive lifesaving interventions than girls have (see figure), a pattern common in South Asian countries. But in the past decade Bangladesh has seen gender disparities effectively disappear in coverage of measles vaccination. As with Brazil, the observed success cannot be attributed to a single initiative, but a series of initiatives aimed at women's empowerment (micro-credit, women's groups, female education and the like) coupled with greater access to health care, particularly through selective outreach by community workers, may account for these changes. An in-depth analysis in the Matlab area shows that community health workers contributed to reducing gender inequities in immunization coverage.³⁹ Nevertheless, socioeconomic disparities in Bangladesh remain large.



Source: Demographic and Health Surveys.



Financing for maternal, newborn and child health




Monitoring financial resource flows for maternal, newborn and child health is a central part of the Countdown-determining the funding gap between resources currently available and the actual investments required to reach national and MDG targets and holding governments and the international community to account for investing adequately in the health of women and children. Policy-makers need financial information to make informed decisions on how to best allocate resources among competing needs, set priorities and ensure sustainable funding for programmes. This section presents an update on the financing gap and patterns in ODA; work on compiling data on national contributions is under way and will be reported in 2011.

Closing the gap: what resources are needed to scale up coverage of maternal, newborn and child interventions?


Preliminary estimates show that if current funding trends continue during 2008–2015, the 68 Countdown countries will face a roughly \$60 billion funding gap relative to the costs of implementing a full package of maternal, newborn and child health Countdown interventions (figure 20). If public commitments by both aid donors and governments are met (assuming a linear increase to 15% of gross national product by 2015 for African countries committed to reaching the Abuja target and 10%–12% for all other *Countdown* countries), the gap remains \$22 billion. This analysis shows that more funding is needed for countries to be able to provide universal coverage of essential maternal, newborn and child health services.

Official development assistance to maternal, newborn and child health shows promising trends

Total ODA for maternal, newborn and child health in 2007 was \$4.1 billion, up 16% from



2006 and nearly double the \$2.1 billion in 2003 (figure 21). Although these trends show improved commitment, ODA for maternal, newborn and child health accounted for only 31% of all ODA for health in 2007, ODA flows for maternal. newborn and child health are important to track, but national resources are a much larger share of funding for maternal, newborn and child health. Even for very low-income Countdown countries such as Ethiopia and Malawi, national sources account for half or more of total spending on reproductive and child health. Tracking government and nongovernment spending at the country level is essential for policy-makers to follow progress in making adequate resources available for women and children. Estimates of domestic expenditure on maternal, newborn and child health, including family planning, will be available in 2011 for a subset of Countdown countries.



Trends in official development assistance by purpose


There are important differences in levels and trends for ODA to child health, maternal and newborn health, and family planning, which is included in Countdown for the first time in this update. Following significant declines in the 1990s, ODA for family planning continued to decline in real terms concurrent with increases in ODA for maternal, newborn and child health (figure 22). This may indicate replacement, with funds targeted to family planning being reallocated to maternal, newborn and child health—especially given that some interventions are delivered in the same service settings. Accurate donor reporting, including correct attribution of funds to specific service areas, is a problem with available data. Better-resourced maternal and child health services result in improvements in maternal, newborn and child survival and other health benefits similar to those that result from strengthened family planning services in many cases. In-depth analyses of these questions are under way and will be reported in 2011.

Is official development assistance targeted to countries in greatest need?

ODA flows are rarely well targeted to either the poorest countries or the countries with the greatest burden of mortality. Table 2 shows

FIGURE 22

Have increases in official development assistance for maternal, newborn and child health substituted for declines (in real terms) in official development assistance for family planning?



Source: Greco and others 2010; UNFPA Donor Support, March 2010, (http://www.unfpa.org/webdav/site/global/shared/documents/publications/2009/2008_donor_support_report.pdf).

TABLE 2

Official development assistance is not always going where it is most needed

Official development assistance to *Countdown* countries, by income group and under-five mortality rate, 2007

GDP per capita (\$)	ODA for maternal, newborn and child health per capita (\$)
Less than 500	0.02-6.40
500-1,000	0.24-6.90
More than 1,000	0.04-28.40
Under-five mortality rate (per 1,000)	ODA for child health per child under age 5 (\$)
More than 200	4.0-28.7
150-200	4.5-225.7
100-149	2.7-18.9
Less than 100	0.3-12.9
Source: Greco and others 2010.	

the distribution of ODA by country income group and under-five mortality level. The wide and overlapping range of ODA for maternal, newborn and child health per capita by country income group and of ODA to child health per child by under-five mortality level indicates the weak link between ODA levels and these two measures of need.

Action now for Millennium Development Goals 4 and 5



The majority of maternal, newborn and child deaths are preventable. Commitments to action are needed by governments and the international community to:

- Make every mother and child count, by registering and counting every birth and death.
- Ensure that every woman, newborn and child has access to essential health services, by overcoming financial barriers.
- Improve equity, by making services available to poor, remote and vulnerable populations.
- Ensure adequate numbers of skilled health workers in every district, by prioritizing training, distribution and retention.

Specific actions for governments and leaders

All countries should:

- Identify inequities in coverage—by geographic area, ethnic group, income and the like—and initiate actions to provide universal coverage of essential interventions and packages.
- Identify gaps in coverage and quality of care along the continuum of care.
- Initiate actions to improve the delivery of essential interventions and packages.
- Increase resource allocations for reproductive, maternal, newborn and child health services, ensuring that interventions and programmes are sufficiently funded.

Parliamentarians should:

- Participate in national and local reviews of health MDG data to monitor progress
- Advocate for greater budgetary resources for maternal, newborn and child health and hold governments to account for meeting promised commitments.
- Review legislative frameworks to be sure evidence based polices for women's and children's health are adopted.

Countries on track to achieving their MDG4 and MDG5 targets should:

- Continue to improve coverage and maintain declines in child, newborn and maternal mortality.
- Ensure that all underserved populations are reached
- Document and share experiences to show how effective policy changes, programme approaches and investments have helped improve maternal, newborn and child health.

Countries making progress towards achieving Millennium Development Goals 4 and 5 should:

- Identify high coverage interventions and document the approaches and investments that supported those achievements.
- Identify low coverage, but high impact, interventions in order to determine how best to provide additional attention and investment.
- Identify and overcome health system constraints and social determinants hindering high coverage.
- Broaden focus to reach underserved populations.

Countries not making progress in reaching Millennium Development Goals 4 and 5 should:

- Identify resource, health system and broader contextual constraints to high coverage.
- Develop national investment and implementation plans for scale-up of interventions proven to reduce maternal, newborn and child mortality.
- Learn from successful local programmes and global research findings
- Adopt and implement evidence-based policies.
- Utilize international expertise and resources.
- Focus on scaling up using innovative strategies.

Countries that have not adopted internationally recommended policies should:

- Review those policies in relation to their own policies and conditions.
- Act on policies that will contribute to improving reproductive, maternal, newborn and child health.

Specific actions for the international community

- Increase and better target donor funding for reproductive, maternal, newborn, and child health through innovative mechanisms and ensure that funding is predictable, consistent and responsive to national needs and plans.
- Support country efforts to improve data collection and analysis by strengthening health information and vital registration systems as well as by undertaking additional surveys to
- measure mortality, coverage and financial flows.
- Invest in implementation research to identify effective strategies for delivering proven interventions and quantify their impact.
- Maximize financial and technical support for large-scale implementation of priority strategies and interventions.
- Encourage the development and use of mechanisms for holding key actors accountable for fulfilling their commitments.



Conclusion



Countdown to 2015: Tracking Progress in Maternal, Newborn and Child Survival was founded on a commitment by academics, governments, international agencies, health care professional associations, donors and nongovernmental organizations to work together towards achieving Millennium Development Goals 4 and 5. By monitoring country progress in the 68 countries that account for more than 95% of all maternal and child deaths and by noting changes in both mortality and coverage of effective interventions, Countdown calls attention to what can be achieved and highlights where countries, interventions and health systems have stalled.

This third *Countdown* report documents changes since 2000, a decade of rapid progress for a few countries and continuing improvements for many others. Despite these positive signs, some

countries have shown little or no improvement in mortality rates and coverage levels or have yet to adopt the policies or evidence-based interventions that save lives. This report explores coverage determinants—including health systems and policies, financial flows and equity—that help explain these differences and acknowledges the importance of social determinants to maternal, newborn and child health and survival.

Countdown partners must work together now to increase their efforts and resources, focusing not just on one intervention or cause but on developing a functional continuum of basic services that save lives and improve health for millions of women, newborns and children. There is still time. This report shows that, by investing our attention and our financial resources, so much more is possible.



Country profiles



These individual country profiles represent the basic information to be analyzed at *Countdown* conferences and evidence for assessing progress since the first Countdown Report in 2005. Each profile presents the most recent available information on selected demographic measures of maternal, newborn and child survival and nutritional status, coverage rates for priority interventions, and selected indicators of equity, policy support, human resources and financial flows.

The information summarized in these pages is intended to help policy-makers and their partners assess progress and prioritize actions in the effort to reduce maternal, newborn and child mortality.

Afghanistan Angola Azerbaijan Bangladesh Benin Bolivia Botswana Brazil

Burkina Faso Burundi Cambodia Cameroon

Central African Republic

Chad China Congo

Congo, Democratic Republic of the

Côte d'Ivoire Djibouti Egypt

Equatorial Guinea

Eritrea Ethiopia Gabon Gambia, The Ghana

Guatemala Guinea

Guinea-Bissau

Haiti India Indonesia Iraq Kenya

Korea, Democratic People's Republic of

Lao People's Democratic Republic

Lesotho Liberia Madagascar Malawi Mali Mauritania Mexico Morocco Mozambique Myanmar Nepal

Nigeria Pakistan

Niger

Papua New Guinea

Peru **Philippines** Rwanda Senegal Sierra Leone Somalia South Africa Sudan Swaziland Tajikistan

Tanzania, United Republic of

Togo


Turkmenistan Uganda Yemen Zambia Zimbabwe




Afghanistan

DEMOGRAPHICS

Total population (000)	27,208	(2008)
Total under-five population (000)	4,907	(2008)
Births (000)	1,269	(2008)
Birth registration (%)	6	(2003)
Under-five mortality rate (per 1000 live births)	257	(2008)
Infant mortality rate (per 1000 live births)	165	(2008)
Neonatal mortality rate (per 1000 live births)	50	(2008)
Total under-five deaths (000)	311	(2008)
Maternal mortality ratio (per 100,000 live births)	1,800	(2005)
Lifetime risk of maternal death (1 in N)	8	(2005)
Total maternal deaths	26,000	(2005)






INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN

Stunting prevalence (moderate and severe, %) 59 (2003-2004) Wasting prevalence (moderate and severe, %) 9 (2003-2004)

Wasting prevalence (moderate and severe, %) 9 (2003-2004)


Underweight prevalence





Complementary feeding rate (6-9 months, %) 29 (2003)


Low birthweight incidence (%) ---



















Adolescent birth rate (births per 1,000 woman)

Unmet need for family planning (%)

Antenatal visits for woman (4 or more visits, %)


Intermittent preventive treatment for malaria (%)

C-section rate (total, urban, rural; %)
(Minimum target is 5% and maximum target is 15%)


Early initiation of breastfeeding (within 1 hr of birth, %)


Postnatal visit for baby (within 2 days for home births, %)


^{*} Not applicable













WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Partial
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	Yes
IMCI adapted to cover newborns 0-1 week of age	No
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	Yes
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	No

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	83 (2007)	
General government expenditure on health as % of total government expenditure (%)	4 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	76 (2007)	
Density of health workers (per 10,000 population)	7.0 (2005)	
Official Development Assistance to child health per child (US\$)	14 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	18 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)		


Afghanistan




Angola

DEMOGRAPHICS

Total population (000)	18,021	(2008)
Total under-five population (000)	3,170	(2008)
Births (000)	774	(2008)
Birth registration (%)	29	(2000)
Under-five mortality rate (per 1000 live births)	220	(2008)
Infant mortality rate (per 1000 live births)	130	(2008)
Neonatal mortality rate (per 1000 live births)	47	(2008)
Total under-five deaths (000)	165	(2008)
Maternal mortality ratio (per 100,000 live births)	1,400	(2005)
Lifetime risk of maternal death (1 in N)	12	(2005)
Total maternal deaths	11,000	(2005)





INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


NUTRITION

Stunting prevalence (moderate and severe, %) 29 (2007) Wasting prevalence (moderate and severe, %) 8 (2007)


107)


Complementary feeding rate (6-9 months, %) Low birthweight incidence (%) 77 (2001)


12 (2000)
























Adolescent birth rate (births per 1,000 women)

Unmet need for family planning (%)

Antenatal visits for woman (4 or more visits, %)


Intermittent preventive treatment for malaria (%)


C-section rate (total, urban, rural; %)
(Minimum target is 5% and maximum target is 15%)


Early initiation of breastfeeding (within 1 hr of birth, %)


Postnatal visit for baby (within 2 days for home births, %)














WATER AND SANITATION





EQUITY



POLICIES

No
Partial
No
No
Partial
Yes
No
Yes

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	131 (2007)	
General government expenditure on health as % of total government expenditure (%)	5 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	20 (2007)	
Density of health workers (per 10,000 population)	14.3 (2004)	
Official Development Assistance to child health per child (US\$)	10 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	11 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)	25 (2006)	


Angola




Azerbaijan

DEMOGRAPHICS

Total population (000)	8,731	(2008)
Total under-five population (000)	738	(2008)
Births (000)	166	(2008)
Birth registration (%)	94	(2006)
Under-five mortality rate (per 1000 live births)	36	(2008)
Infant mortality rate (per 1000 live births)	32	(2008)
Neonatal mortality rate (per 1000 live births)	19	(2008)
Total under-five deaths (000)	6	(2008)
Maternal mortality ratio (per 100,000 live births)	82	(2005)
Lifetime risk of maternal death (1 in N)	670	(2005)
Total maternal deaths	110	(2005)





INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN

Stunting prevalence (moderate and severe, %) 25 (2006) Wasting prevalence (moderate and severe, %)


7 (2006)


Low birthweight incidence (%)


Complementary feeding rate (6-9 months, %)


44 (2006)


10 (2006)
























Adolescent birth rate (births per 1.000 women) 42 (2008) Unmet need for family planning (%) 23 (2006)Antenatal visits for woman (4 or more visits, %) 45 (2006) Intermittent preventive treatment for malaria (%) NA* C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) 5, 5, **4** (2006)Early initiation of breastfeeding (within 1 hr of birth, %) 32 (2006)


Postnatal visit for baby (within 2 days for home births, %) ---

















WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Partial
New ORS formula and zinc for management of diarrhoea	No
Community treatment of pneumonia with antibiotics	Partial
IMCI adapted to cover newborns 0-1 week of age	No
Costed implementation plan(s) for maternal, newborn and child health available	No
Midwives be authorised to administer a core set of life saving interventions	Partial
Maternity protection in accordance with ILO Convention 183	Partial
Specific notification of maternal deaths	Yes

SYSTEMS

Financial Flows and Human Resources	
Per capita total expenditure on health (US\$)	284 (2007)
General government expenditure on health as % of total government expenditure (%)	4 (2007)
Out-of-pocket expenditure as % of total expenditure on health (%)	64 (2007)
Density of health workers (per 10,000 population)	122.1 (2007)
Official Development Assistance to child health per child (US\$)	2 (2007)
Official Development Assistance to maternal an neonatal health per live birth (US\$)	d 8 (2007)
National availability of Emergency Obstetric Car services (% of recommended minimum)	re


Azerbaijan




Bangladesh

DEMOGRAPHICS

Total population (000)	160,000	(2008)
Total under-five population (000)	16,710	(2008)
Births (000)	3,430	(2008)
Birth registration (%)	10	(2006)
Under-five mortality rate (per 1000 live births)	54	(2008)
Infant mortality rate (per 1000 live births)	43	(2008)
Neonatal mortality rate (per 1000 live births)	33	(2008)
Total under-five deaths (000)	183	(2008)
Maternal mortality ratio (per 100,000 live births	570	(2005)
Lifetime risk of maternal death (1 in N)	51	(2005)
Total maternal deaths	21,000	(2005)






INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %) 43 (2007) Wasting prevalence (moderate and severe, %)


17 (2007)


Complementary feeding rate (6-9 months, %) Low birthweight incidence (%)


74 (2007) 22 (2006)























Adolescent birth rate (births per 1,000 women)

Unmet need for family planning (%)

Antenatal visits for woman (4 or more visits, %)

Intermittent preventive treatment for malaria (%)


C-section rate (total, urban, rural; %)

C-section rate (total, urban, **rural**; %) (Minimum target is 5% and maximum target is 15%) 8, 16, **5** (2007)


Early initiation of breastfeeding (within 1 hr of birth, %)


Postnatal visit for baby (within 2 days for home births, %) 19 (2007)


43 (2007)














WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Partial
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	Yes
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Partial
Midwives be authorised to administer a core set of life saving interventions	Partial
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	Partial

SYSTEMS

Financial Flows and Human Reso	ources
Per capita total expenditure on health (US\$)	42 (2007)
General government expenditure on health as % of total government expenditure (%)	8 (2007)
Out-of-pocket expenditure as % of total expenditure on health (%)	65 (2007)
Density of health workers (per 10,000 population)	5.8 (2005)
Official Development Assistance to child health per child (US\$)	3 (2007)
Official Development Assistance to maternal and neonatal health per live birth (US\$)	8 (2007)
National availability of Emergency Obstetric Care services (% of recommended minimum)	


Bangladesh




Benin

DEMOGRAPHICS

Total population (000)	8,662	(2008)
Total under-five population (000)	1,450	(2008)
Births (000)	342	(2008)
Birth registration (%)	60	(2006)
Under-five mortality rate (per 1000 live births)	121	(2008)
Infant mortality rate (per 1000 live births)	76	(2008)
Neonatal mortality rate (per 1000 live births)	33	(2008)
Total under-five deaths (000)	39	(2008)
Maternal mortality ratio (per 100,000 live births)	840	(2005)
Lifetime risk of maternal death (1 in N)	20	(2005)
Total maternal deaths	2,900	(2005)






INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Malaria prevention


Percent children < 5 years sleeping under ITNs


Stunting prevalence (moderate and severe, %) Wasting prevalence (moderate and severe, %)


43 (2006) 8 (2006) Complementary feeding rate (6-9 months, %) Low birthweight incidence (%) 72 (2006) 15 (2006)



























Adolescent birth rate (births per 1,000 women) 114 (2004)
Unmet need for family planning (%) 30 (2006)
Antenatal visits for woman (4 or more visits, %) 61 (2006)
Intermittent preventive treatment for malaria (%) 3 (2006)
C-section rate (total, urban, rural; %)
(Minimum target is 5% and maximum target is 15%) 4, 6, 2 (2006)
Early initiation of breastfeeding (within 1 hr of birth, %) 54 (2006)
Postnatal visit for baby (within 2 days for home births, %) ---














WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Yes
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	Partial
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	Partial
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	Yes

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	70 (2007)	
General government expenditure on health as % of total government expenditure (%)	11 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	46 (2007)	
Density of health workers (per 10,000 population)	8.3 (2008)	
Official Development Assistance to child health per child (US\$)	12 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	14 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)	66 (2002)	


Benin




Bolivia

DEMOGRAPHICS

Total population (000)	9,694	(2008)
Total under-five population (000)	1,245	(2008)
Births (000)	263	(2008)
Birth registration (%)	74	(2001)
Under-five mortality rate (per 1000 live births)	54	(2008)
Infant mortality rate (per 1000 live births)	46	(2008)
Neonatal mortality rate (per 1000 live births)	25	(2008)
Total under-five deaths (000)	14	(2008)
Maternal mortality ratio (per 100,000 live births)	290	(2005)
Lifetime risk of maternal death (1 in N)	89	(2005)
Total maternal deaths	760	(2005)







INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


NUTRITION


Stunting prevalence (moderate and severe, %) 32 (2003) Wasting prevalence (moderate and severe, %) 2 (2003)

Complementary feeding rate (6-9 months, %) Low birthweight incidence (%) 81 (2008) 7 (2003)
























Adolescent birth rate (births per 1,000 women) 89 (2006)
Unmet need for family planning (%) 20 (2008)
Antenatal visits for woman (4 or more visits, %) 58 (2003)
Intermittent preventive treatment for malaria (%) NA*


C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) 15,21,6 (2003)

Early initiation of breastfeeding (within 1 hr of birth, %) 61 (2003)


Postnatal visit for baby (within 2 days for home births, %)

















WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Partial
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	Partial
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	Partial
Maternity protection in accordance with ILO Convention 183	Partial
Specific notification of maternal deaths	Partial

SYSTEMS

Financial Flows and Human Res	ources
Per capita total expenditure on health (US\$)	219 (2007)
General government expenditure on health as % of total government expenditure (%)	10 (2007)
Out-of-pocket expenditure as % of total expenditure on health (%)	32 (2007)
Density of health workers (per 10,000 population)	33.5 (2001)
Official Development Assistance to child health per child (US\$)	14 (2007)
Official Development Assistance to maternal and neonatal health per live birth (US\$)	20 (2007)
National availability of Emergency Obstetric Care services (% of recommended minimum)	48 (2003)


Bolivia




Botswana

DEMOGRAPHICS

Total population (000)	1,921	(2008)
Total under-five population (000)	221	(2008)
Births (000)	47	(2008)
Birth registration (%)	58	(2000)
Under-five mortality rate (per 1000 live births)	31	(2008)
Infant mortality rate (per 1000 live births)	26	(2008)
Neonatal mortality rate (per 1000 live births)	16	(2008)
Total under-five deaths (000)	1	(2008)
Maternal mortality ratio (per 100,000 live births)	380	(2005)
Lifetime risk of maternal death (1 in N)	130	(2005)
Total maternal deaths	170	(2005)







INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


NUTRITION


Stunting prevalence (moderate and severe, %) 29 (2000) Wasting prevalence (moderate and severe, %) 6 (2000)


Complementary feeding rate (6-9 months, %) Low birthweight incidence (%) 57 (2000) 10 (2000)























Adolescent birth rate (births per 1,000 women) 51 (2006)

Unmet need for family planning (%) 27 (1998)


Antenatal visits for woman (4 or more visits, %) ---
Intermittent preventive treatment for malaria (%) NA*


C-section rate (total, urban, rural; %)
(Minimum target is 5% and maximum target is 15%) ----, ----, ---
Early initiation of breastfeeding (within 1 hr of birth, %) ---
Postnatal visit for baby (within 2 days for home births, %) ----
















WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Yes
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	No
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	Partial
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	Yes

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	762 (2007)	
General government expenditure on health as % of total government expenditure (%)	13 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	7 (2007)	
Density of health workers (per 10,000 population)	30.5 (2004)	
Official Development Assistance to child health per child (US\$)	3 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	16 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)		


Botswana




Brazil

DEMOGRAPHICS

Total population (000)	191,972	(2008)
Total under-five population (000)	16,125	(2008)
Births (000)	3,105	(2008)
Birth registration (%)	89	(2005)
Under-five mortality rate (per 1000 live births)	22	(2008)
Infant mortality rate (per 1000 live births)	18	(2008)
Neonatal mortality rate (per 1000 live births)	11	(2008)
Total under-five deaths (000)	67	(2008)
Maternal mortality ratio (per 100,000 live births))	
Lifetime risk of maternal death (1 in N)		
Total maternal deaths	4,100	(2005)






INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %) 7 (2006) Wasting prevalence (moderate and severe, %)


2 (2006)


Complementary feeding rate (6-9 months, %) Low birthweight incidence (%)


70 (2006) 8 (2006)
























Adolescent birth rate (births per 1,000 women) 56 (2006)
Unmet need for family planning (%) 7 (1996)
Antenatal visits for woman (4 or more visits, %) 89 (2006)
Intermittent preventive treatment for malaria (%) NA*


C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) 46, 35, **45** (2006) Early initiation of breastfeeding (within 1 hr of birth, %) 43 (2006)


Postnatal visit for baby (within 2 days for home births, %)














WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Yes
New ORS formula and zinc for management of diarrhoea	No
Community treatment of pneumonia with antibiotics	No
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Partial
Midwives be authorised to administer a core set of life saving interventions	Partial
Maternity protection in accordance with ILO Convention 183	Partial
Specific notification of maternal deaths	Yes

SYSTEMS

Financial Flows and Human Resources			
Per capita total expenditure on health (US\$)	837 (2007)		
General government expenditure on health as % of total government expenditure (%)	5 (2007)		
Out-of-pocket expenditure as % of total expenditure on health (%)	34 (2007)		
Density of health workers (per 10,000 population)	46.0 (2006)		
Official Development Assistance to child health per child (US\$)	0 (2007)		
Official Development Assistance to maternal and neonatal health per live birth (US\$)	0 (2007)		
National availability of Emergency Obstetric Care services (% of recommended minimum)			


Brazil




Burkina Faso

DEMOGRAPHICS

Total population (000)	15,234	(2008)
Total under-five population (000)	2,934	(2008)
Births (000)	721	(2008)
Birth registration (%)	64	(2006)
Under-five mortality rate (per 1000 live births)	169	(2008)
Infant mortality rate (per 1000 live births)	92	(2008)
Neonatal mortality rate (per 1000 live births)	36	(2008)
Total under-five deaths (000)	117	(2008)
Maternal mortality ratio (per 100,000 live births)	700	(2005)
Lifetime risk of maternal death (1 in N)	22	(2005)
Total maternal deaths	4,300	(2005)






INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %) 41 (2006) Wasting prevalence (moderate and severe, %)


25 (2006)


Complementary feeding rate (6-9 months, %) Low birthweight incidence (%)


50 (2006) 16 (2006)























Adolescent birth rate (births per 1,000 women) 131 (2001)

Unmet need for family planning (%) 29 (2003)

Antenatal visits for woman (4 or more visits, %) 18 (2003)


Intermittent preventive treatment for malaria (%) 1 (2006)


C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) 1, 3, 0 (2003)


Early initiation of breastfeeding (within 1 hr of birth, %) 20 (2006)


Postnatal visit for baby (within 2 days for home births, %) ---













WATER AND SANITATION







Mean coverage

3rd

Wealth quintile

Wealthiest

EQUITY

20

0

Source: DHS 2003

Poorest

POLICIES

International Code of Marketing of Breastmilk Substitutes	Yes
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	No
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	Yes
Maternity protection in accordance with ILO Convention 183	Partial
Specific notification of maternal deaths	Yes

SYSTEMS

Financial Flows and Human Resources			
Per capita total expenditure on health (US\$)	72 (2007)		
General government expenditure on health as % of total government expenditure (%)	13 (2007)		
Out-of-pocket expenditure as % of total expenditure on health (%)	40 (2007)		
Density of health workers (per 10,000 population)	7.9 (2008)		
Official Development Assistance to child health per child (US\$)	10 (2007)		
Official Development Assistance to maternal and neonatal health per live birth (US\$)	14 (2007)		
National availability of Emergency Obstetric Care services (% of recommended minimum)			


Burkina Faso




Burundi

DEMOGRAPHICS

Total population (000)	8,074	(2008)
Total under-five population (000)	1,155	(2008)
Births (000)	278	(2008)
Birth registration (%)	60	(2005)
Under-five mortality rate (per 1000 live births)	168	(2008)
Infant mortality rate (per 1000 live births)	102	(2008)
Neonatal mortality rate (per 1000 live births)	42	(2008)
Total under-five deaths (000)	45	(2008)
Maternal mortality ratio (per 100,000 live births)	1,100	(2005)
Lifetime risk of maternal death (1 in N)	16	(2005)
Total maternal deaths	3,900	(2005)







INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


NUTRITION


Stunting prevalence (moderate and severe, %) 63 (2005)
Wasting prevalence (moderate and severe, %) 8 (2005)


Complementary feeding rate (6-9 months, %) Low birthweight incidence (%) 88 (2005) 11 (2005)
























Adolescent birth rate (births per 1,000 women) 30 (2001)

Unmet need for family planning (%) 29 (2002)


Antenatal visits for woman (4 or more visits, %) --
Intermittent preventive treatment for malaria (%) NA*


C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) ---, ---, --
Early initiation of breastfeeding (within 1 hr of birth, %) --
Postnatal visit for baby (within 2 days for home births, %) ---


^{*} Not applicable














WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Partial
New ORS formula and zinc for management of diarrhoea	Partial
Community treatment of pneumonia with antibiotics	No
IMCI adapted to cover newborns 0-1 week of age	Partial
Costed implementation plan(s) for maternal, newborn and child health available	Partial
Midwives be authorised to administer a core set of life saving interventions	No
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	Partial

SYSTEMS

Financial Flows and Human Resources			
Per capita total expenditure on health (US\$)	51 (2007)		
General government expenditure on health as % of total government expenditure (%)	13 (2007)		
Out-of-pocket expenditure as % of total expenditure on health (%)	38 (2007)		
Density of health workers (per 10,000 population)	2.2 (2004)		
Official Development Assistance to child health per child (US\$)	15 (2007)		
Official Development Assistance to maternal and neonatal health per live birth (US\$)	19 (2007)		
National availability of Emergency Obstetric Care services (% of recommended minimum)			

Burundi




Cambodia

DEMOGRAPHICS

Total population (000)	14,562	(2008)
Total under-five population (000)	1,611	(2008)
Births (000)	361	(2008)
Birth registration (%)	66	(2005)
Under-five mortality rate (per 1000 live births)	90	(2008)
Infant mortality rate (per 1000 live births)	69	(2008)
Neonatal mortality rate (per 1000 live births)	31	(2008)
Total under-five deaths (000)	32	(2008)
Maternal mortality ratio (per 100,000 live births)	540	(2005)
Lifetime risk of maternal death (1 in N)	48	(2005)
Total maternal deaths	2,300	(2005)







INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


NUTRITION


Stunting prevalence (moderate and severe, %) 42 (2005)
Wasting prevalence (moderate and severe, %) 9 (2005)


Complementary feeding rate (6-9 months, %) Low birthweight incidence (%) 82 (2005) 14 (2005)























Adolescent birth rate (births per 1,000 women) 52 (2003)

Unmet need for family planning (%) 25 (2005)

Antenatal visits for woman (4 or more visits, %) 27 (2005)

Intermittent preventive treatment for malaria (%) NA*


C-section rate (total, urban, rural; %)
(Minimum target is 5% and maximum target is 15%)

2, 6, 1 (2005)


Early initiation of breastfeeding (within 1 hr of birth, %) 35 (2005)


Postnatal visit for baby (within 2 days for home births, %) ---


* Not applicable














WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Partial
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	Yes
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	Partial
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	Partial

SYSTEMS

Financial Flows and Human Resources			
Per capita total expenditure on health (US\$)	108 (2007)		
General government expenditure on health as % of total government expenditure (%)	11 (2007)		
Out-of-pocket expenditure as % of total expenditure on health (%)	60 (2007)		
Density of health workers (per 10,000 population)	10.1 (2000)		
Official Development Assistance to child health per child (US\$)	8 (2007)		
Official Development Assistance to maternal and neonatal health per live birth (US\$)	27 (2007)		
National availability of Emergency Obstetric Care services (% of recommended minimum)	33 (2008)		


Cambodia




Cameroon

DEMOGRAPHICS

Total population (000)	19,088	(2008)
Total under-five population (000)	3,016	(2008)
Births (000)	704	(2008)
Birth registration (%)	70	(2006)
Under-five mortality rate (per 1000 live births)	131	(2008)
Infant mortality rate (per 1000 live births)	82	(2008)
Neonatal mortality rate (per 1000 live births)	31	(2008)
Total under-five deaths (000)	89	(2008)
Maternal mortality ratio (per 100,000 live births)	1,000	(2005)
Lifetime risk of maternal death (1 in N)	24	(2005)
Total maternal deaths	5,700	(2005)





INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %) 36 (2006) Wasting prevalence (moderate and severe, %)


7 (2006)

Complementary feeding rate (6-9 months, %) Low birthweight incidence (%)


64 (2006)


11 (2006)






























Adolescent birth rate (births per 1,000 women) 141 (2002)
Unmet need for family planning (%) 20 (2004)
Antenatal visits for woman (4 or more visits, %) 60 (2004)
Intermittent preventive treatment for malaria (%) 6 (2006)
C-section rate (total, urban, rural; %)
(Minimum target is 5% and maximum target is 15%) 2, 4, 1 (2004)
Early initiation of breastfeeding (within 1 hr of birth, %) 20 (2006)
Postnatal visit for baby (within 2 days for home births, %) ---














WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Yes
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	Yes
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	Yes
Maternity protection in accordance with ILO Convention 183	Partial
Specific notification of maternal deaths	No

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	104 (2007)	
General government expenditure on health as % of total government expenditure (%)	8 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	70 (2007)	
Density of health workers (per 10,000 population)	17.9 (2004)	
Official Development Assistance to child health per child (US\$)	4 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	6 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)	29 (2000)	


Cameroon




Central African Republic

DEMOGRAPHICS

Total population (000)	4,339	(2008)
Total under-five population (000)	656	(2008)
Births (000)	154	(2008)
Birth registration (%)	49	(2006)
Under-five mortality rate (per 1000 live births)	173	(2008)
Infant mortality rate (per 1000 live births)	115	(2008)
Neonatal mortality rate (per 1000 live births)	47	(2008)
Total under-five deaths (000)	26	(2008)
Maternal mortality ratio (per 100,000 live births)	980	(2005)
Lifetime risk of maternal death (1 in N)	25	(2005)
Total maternal deaths	1,500	(2005)






INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %) 43 (2006) Wasting prevalence (moderate and severe, %)


12 (2006)


Complementary feeding rate (6-9 months, %) Low birthweight incidence (%)


55 (2006) 13 (2006)























Adolescent birth rate (births per 1,000 women) 133 (2003)

Unmet need for family planning (%) 16 (1994-95)


Antenatal visits for woman (4 or more visits, %) 40 (1994-95)

Intermittent preventive treatment for malaria (%) 9 (2006)


C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) 2, 2, 2 (1994-95)


Early initiation of breastfeeding (within 1 hr of birth, %) 39 (2006)


Postnatal visit for baby (within 2 days for home births, %) ---













WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	No
New ORS formula and zinc for management of diarrhoea	No
Community treatment of pneumonia with antibiotics	No
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	Yes
Maternity protection in accordance with ILO Convention 183	Partial
Specific notification of maternal deaths	No

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	30 (2007)	
General government expenditure on health as % of total government expenditure (%)	11 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	62 (2007)	
Density of health workers (per 10,000 population)	4.9 (2004)	
Official Development Assistance to child health per child (US\$)	10 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	23 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)		

Central African Republic




Chad

DEMOGRAPHICS

Total population (000)	10,914	(2008)
Total under-five population (000)	1,985	(2008)
Births (000)	498	(2008)
Birth registration (%)	9	(2004)
Under-five mortality rate (per 1000 live births)	209	(2008)
Infant mortality rate (per 1000 live births)	124	(2008)
Neonatal mortality rate (per 1000 live births)	44	(2008)
Total under-five deaths (000)	99	(2008)
Maternal mortality ratio (per 100,000 live births)	1,500	(2005)
Lifetime risk of maternal death (1 in N)	11	(2005)
Total maternal deaths	6,900	(2005)







INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %)


Wasting prevalence (moderate and severe, %)





Complementary feeding rate (6-9 months, %) 77 (2004) Low birthweight incidence (%) 22 (2004)




















Adolescent birth rate (births per 1,000 woman) 193 (2002)

Unmet need for family planning (%) 23 (2004)


Antenatal visits for woman (4 or more visits, %) 18 (2004)


Intermittent preventive treatment for malaria (%) ---
C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) 0, 1, 0 (2004)

Early initiation of breastfeeding (within 1 hr of birth, %) 34 (2004)


Postnatal visit for baby (within 2 days for home births, %) ---














WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	No
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	No
IMCI adapted to cover newborns 0-1 week of age	No
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	Yes
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	No

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	72 (2007)	
General government expenditure on health as % of total government expenditure (%)	14 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	42 (2007)	
Density of health workers (per 10,000 population)	3.2 (2004)	
Official Development Assistance to child health per child (US\$)	6 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	8 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)	40 (2002)	


Chad




China

DEMOGRAPHICS

Total population (000)	1,337,411	(2008)
Total under-five population (000)	86,881	(2008)
Births (000)	18,134	(2008)
Birth registration (%)		
Under-five mortality rate (per 1000 live birth	ns) 21	(2008)
Infant mortality rate (per 1000 live births)	18	(2008)
Neonatal mortality rate (per 1000 live births	s) 11	(2008)
Total under-five deaths (000)	365	(2008)
Maternal mortality ratio (per 100,000 live bird	ths) 45	(2005)
Lifetime risk of maternal death (1 in N)	1,300	(2005)
Total maternal deaths	7,800	(2005)





INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN

Stunting prevalence (moderate and severe, %) 15 (2005)

Wasting prevalence (moderate and severe, %)


6

2005

Underweight prevalence Percent children < 5 years underweight for age* 100 80 60

Complementary feeding rate (6-9 months, %) 32 (2003) Low birthweight incidence (%) 4 (1999-2003)





ised on 2006 WHO reference population **CHILD HEALTH**

2002


Other NS

Diarrhoeal disease treatment

therapy or increased fluids, with continued feeding


40 20

0





No data














Adolescent birth rate (births per 1,000 women) 5 (2006)
Unmet need for family planning (%) 2 (2001)
Antenatal visits for woman (4 or more visits, %) --Intermittent preventive treatment for malaria (%) NA*
C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) ---, ---, --Early initiation of breastfeeding (within 1 hr of birth, %) --Postnatal visit for baby (within 2 days for home births, %) ---


^{*} Not applicable













WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Partial
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	Yes
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	Partial
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	Yes

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	233 (2007)	
General government expenditure on health as % of total government expenditure (%)	10 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	51 (2007)	
Density of health workers (per 10,000 population)	23.8 (2003)	
Official Development Assistance to child health per child (US\$)	0 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	1 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)		


China




Congo

DEMOGRAPHICS

Total population (000)	3,615	(2008)
Total under-five population (000)	551	(2008)
Births (000)	125	(2008)
Birth registration (%)	81	(2005)
Under-five mortality rate (per 1000 live births)	127	(2008)
Infant mortality rate (per 1000 live births)	80	(2008)
Neonatal mortality rate (per 1000 live births)	34	(2008)
Total under-five deaths (000)	16	(2008)
Maternal mortality ratio (per 100,000 live births)	740	(2005)
Lifetime risk of maternal death (1 in N)	22	(2005)
Total maternal deaths	1,300	(2005)






INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %) 30 (2005) Wasting prevalence (moderate and severe, %)


8 (2005)


Complementary feeding rate (6-9 months, %) Low birthweight incidence (%)


78 (2005) 13 (2005)























Adolescent birth rate (births per 1,000 women) 132 (2003)

Unmet need for family planning (%) 16 (2005)


Antenatal visits for woman (4 or more visits, %) 75 (2005)


Intermittent preventive treatment for malaria (%) ---
C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) 3, 4, 2 (2005)


Early initiation of breastfeeding (within 1 hr of birth, %) 39 (2005)


Postnatal visit for baby (within 2 days for home births, %) ---














WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	No
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	No
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	Yes
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	No

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	90 (2007)	
General government expenditure on health as % of total government expenditure (%)	5 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	30 (2007)	
Density of health workers (per 10,000 population)	9.2 (2007)	
Official Development Assistance to child health per child (US\$)	7 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	8 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)		


Congo




Congo, Democratic Republic of the

DEMOGRAPHICS

Total population (000)	64,257	(2008)
Total under-five population (000)	11,829	(2008)
Births (000)	2,886	(2008)
Birth registration (%)	31	(2007)
Under-five mortality rate (per 1000 live births)	199	(2008)
Infant mortality rate (per 1000 live births)	126	(2008)
Neonatal mortality rate (per 1000 live births)	56	(2008)
Total under-five deaths (000)	554	(2008)
Maternal mortality ratio (per 100,000 live births)	1,100	(2005)
Lifetime risk of maternal death (1 in N)	13	(2005)
Total maternal deaths	32,000	(2005)





INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN

Stunting prevalence (moderate and severe, %) 46 (2007) Wasting prevalence (moderate and severe, %)


10 (2007)


Low birthweight incidence (%)


Complementary feeding rate (6-9 months, %)


82 (2007)


12 (2001)



























Adolescent birth rate (births per 1,000 women) 127 (2005)
Unmet need for family planning (%) 24 (2007)
Antenatal visits for woman (4 or more visits, %) 47 (2007)
Intermittent preventive treatment for malaria (%) 5 (2007)
C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) 4, 5, 4 (2007)
Early initiation of breastfeeding (within 1 hr of birth, %) 48 (2007)
Postnatal visit for baby (within 2 days for home births, %) ---














WATER AND SANITATION





EQUITY



POLICIES

Partial
Yes
No
Yes

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	17 (2007)	
General government expenditure on health as % of total government expenditure (%)	6 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	41 (2007)	
Density of health workers (per 10,000 population)	6.4 (2004)	
Official Development Assistance to child health per child (US\$)	5 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	6 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)		


Congo, Democratic Republic of the




Côte d'Ivoire

DEMOGRAPHICS

Total population (000)	20,591	(2008)
Total under-five population (000)	3,139	(2008)
Births (000)	722	(2008)
Birth registration (%)	55	(2006)
Under-five mortality rate (per 1000 live births)	114	(2008)
Infant mortality rate (per 1000 live births)	81	(2008)
Neonatal mortality rate (per 1000 live births)	41	(2008)
Total under-five deaths (000)	79	(2008)
Maternal mortality ratio (per 100,000 live births)	810	(2005)
Lifetime risk of maternal death (1 in N)	27	(2005)
Total maternal deaths	5,400	(2005)





INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN

Stunting prevalence (moderate and severe, %) 40 (2006) Wasting prevalence (moderate and severe, %)


8 (2006)

Complementary feeding rate (6-9 months, %) Low birthweight incidence (%)


54 (2006)


17 (2006)


Underweight prevalence Percent children < 5 years underweight for age* 100 80 60



























Adolescent birth rate (births per 1,000 women) 111 (2006)Unmet need for family planning (%) 28 (1998-99) Antenatal visits for woman (4 or more visits, %) (2005) Intermittent preventive treatment for malaria (%) 8 (2006)C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) 6, 8, **6** (2005)Early initiation of breastfeeding (within 1 hr of birth, %) 25 (2006) Postnatal visit for baby (within 2 days for home births, %)














WATER AND SANITATION







3rd

Wealth quintile

Wealthiest

EQUITY

0

Source: MICS 2006

Poorest

POLICIES

International Code of Marketing of Breastmilk Substitutes	No
New ORS formula and zinc for management of diarrhoea	Partial
Community treatment of pneumonia with antibiotics	Partial
IMCI adapted to cover newborns 0-1 week of age	No
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	Partial
Maternity protection in accordance with ILO Convention 183	Partial
Specific notification of maternal deaths	Partial

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	67 (2007)	
General government expenditure on health as % of total government expenditure (%)	5 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	67 (2007)	
Density of health workers (per 10,000 population)	6.2 (2008)	
Official Development Assistance to child health per child (US\$)	5 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	7 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)		


Côte d'Ivoire




Djibouti

DEMOGRAPHICS

Total population (000)	849	(2008)
Total under-five population (000)	108	(2008)
Births (000)	24	(2008)
Birth registration (%)	89	(2006)
Under-five mortality rate (per 1000 live births)	95	(2008)
Infant mortality rate (per 1000 live births)	76	(2008)
Neonatal mortality rate (per 1000 live births)	36	(2008)
Total under-five deaths (000)	2	(2008)
Maternal mortality ratio (per 100,000 live births)	650	(2005)
Lifetime risk of maternal death (1 in N)	35	(2005)
Total maternal deaths	180	(2005)






INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %) 33 (2007) Wasting prevalence (moderate and severe, %) 17 (2007)


Complementary feeding rate (6-9 months, %) Low birthweight incidence (%)


23 (2006)


10 (2006)























Adolescent birth rate (births per 1,000 women)

Unmet need for family planning (%)

Antenatal visits for woman (4 or more visits, %)

7 (2002)

Intermittent preventive treatment for malaria (%)


C-section rate (total, urban, rural; %)
(Minimum target is 5% and maximum target is 15%)


Early initiation of breastfeeding (within 1 hr of birth, %)


55 (2006)

Postnatal visit for baby (within 2 days for home births, %)














WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Partial
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	No
IMCI adapted to cover newborns 0-1 week of age	No
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	Partial
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	Yes

SYSTEMS


Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	148 (2007)	
General government expenditure on health as % of total government expenditure (%)	14 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	23 (2007)	
Density of health workers (per 10,000 population)	7.5 (2005)	
Official Development Assistance to child health per child (US\$)	66 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	108 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)	50 (2004)	


Djibouti



DEMOGRAPHICS

Total population (000)	81,527	(2008)
Total under-five population (000)	9,447	(2008)
Births (000)	2,015	(2008)
Birth registration (%)	99	(2005)
Under-five mortality rate (per 1000 live births)	23	(2008)
Infant mortality rate (per 1000 live births)	20	(2008)
Neonatal mortality rate (per 1000 live births)	13	(2008)
Total under-five deaths (000)	45	(2008)
Maternal mortality ratio (per 100,000 live births)	130	(2005)
Lifetime risk of maternal death (1 in N)	230	(2005)
Total maternal deaths	2,400	(2005)





INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN

Malaria prevention

Percent children < 5 years sleeping under ITNs*

Stunting prevalence (moderate and severe, %)


Wasting prevalence (moderate and severe, %)

29 (2008)


7 (2008)


Complementary feeding rate (6-9 months, %) Low birthweight incidence (%)

66 (2008) 13 (2008)






















Adolescent birth rate (births per 1,000 women) 50 (2006)
Unmet need for family planning (%) 9 (2008)
Antenatal visits for woman (4 or more visits, %) 66 (2008)
Intermittent preventive treatment for malaria (%) NA*


C-section rate (total, urban, rural; %)

 $(Minimum\ target\ is\ 5\%\ and\ maximum\ target\ is\ 15\%) \qquad 28,37, \textcolor{red}{\textbf{22}}\ \ (2008)$


Early initiation of breastfeeding (within 1 hr of birth, %) 56 (2008)


Postnatal visit for baby (within 2 days for home births, %) 8 (2008)


* Not applicable














WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Partial
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	No
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	Partial
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	Yes

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	310 (2007)	
General government expenditure on health as % of total government expenditure (%)	7 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	59 (2007)	
Density of health workers (per 10,000 population)	57.8 (2005)	
Official Development Assistance to child health per child (US\$)	2 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	4 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)		


Egypt




Equatorial Guinea

DEMOGRAPHICS

Total population (000)	659	(2008)
Total under-five population (000)	103	(2008)
Births (000)	25	(2008)
Birth registration (%)	32	(2000)
Under-five mortality rate (per 1000 live births)	148	(2008)
Infant mortality rate (per 1000 live births)	90	(2008)
Neonatal mortality rate (per 1000 live births)	40	(2008)
Total under-five deaths (000)	3	(2008)
Maternal mortality ratio (per 100,000 live births)	680	(2005)
Lifetime risk of maternal death (1 in N)	28	(2005)
Total maternal deaths	150	(2005)





INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN

100

80

60

40

20 0

Complementary feeding rate (6-9 months, %)

Percent infants < 6 months exclusively breastfed

Exclusive breastfeeding

Stunting prevalence (moderate and severe, %)

Wasting prevalence (moderate and severe, %)

Low birthweight incidence (%) 13 (2000)


24


2000





No data























Adolescent birth rate (births per 1,000 women)

Unmet need for family planning (%)

Antenatal visits for woman (4 or more visits, %)


Intermittent preventive treatment for malaria (%)


C-section rate (total, urban, rural; %)
(Minimum target is 5% and maximum target is 15%)


Early initiation of breastfeeding (within 1 hr of birth, %)


Postnatal visit for baby (within 2 days for home births, %)













WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	No
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	Yes
IMCI adapted to cover newborns 0-1 week of age	No
Costed implementation plan(s) for maternal, newborn and child health available	Partial
Midwives be authorised to administer a core set of life saving interventions	Partial
Maternity protection in accordance with ILO Convention 183	Partial
Specific notification of maternal deaths	Partial

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	543 (2007)	
General government expenditure on health as % of total government expenditure (%)	7 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	15 (2007)	
Density of health workers (per 10,000 population)	8.3 (2004)	
Official Development Assistance to child health per child (US\$)	22 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	52 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)		


Equatorial Guinea




Eritrea

DEMOGRAPHICS

Total population (000)	4,927	(2008)
Total under-five population (000)	811	(2008)
Births (000)	182	(2008)
Birth registration (%)		
Under-five mortality rate (per 1000 live births)	58	(2008)
Infant mortality rate (per 1000 live births)	41	(2008)
Neonatal mortality rate (per 1000 live births)	17	(2008)
Total under-five deaths (000)	10	(2008)
Maternal mortality ratio (per 100,000 live births)	450	(2005)
Lifetime risk of maternal death (1 in N)	44	(2005)
Total maternal deaths	760	(2005)







INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


NUTRITION


Stunting prevalence (moderate and severe, %) 44 (2002) Wasting prevalence (moderate and severe, %) 15 (2002)


Complementary feeding rate (6-9 months, %) Low birthweight incidence (%) 43 (2002) 14 (2002)























Adolescent birth rate (births per 1,000 women)

Unmet need for family planning (%)

Antenatal visits for woman (4 or more visits, %)

Intermittent preventive treatment for malaria (%)


NA*

C-section rate (total, urban, **rural**; %) (Minimum target is 5% and maximum target is 15%) 3, 7, 1 (2002)


Early initiation of breastfeeding (within 1 hr of birth, %) 78 (2002)


Postnatal visit for baby (within 2 days for home births, %) ---


* Not applicable














WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	No
New ORS formula and zinc for management of diarrhoea	Partial
Community treatment of pneumonia with antibiotics	Partial
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Partial
Midwives be authorised to administer a core set of life saving interventions	Partial
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	No

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	20 (2007)	
General government expenditure on health as % of total government expenditure (%)	4 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	55 (2007)	
Density of health workers (per 10,000 population)	6.3 (2004)	
Official Development Assistance to child health per child (US\$)	9 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	13 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)		


Eritrea




Ethiopia

DEMOGRAPHICS

Total population (000)	80,713	(2008)
Total under-five population (000)	13,323	(2008)
Births (000)	3,093	(2008)
Birth registration (%)	7	(2005)
Under-five mortality rate (per 1000 live births)	109	(2008)
Infant mortality rate (per 1000 live births)	69	(2008)
Neonatal mortality rate (per 1000 live births)	39	(2008)
Total under-five deaths (000)	321	(2008)
Maternal mortality ratio (per 100,000 live births)	720	(2005)
Lifetime risk of maternal death (1 in N)	27	(2005)
Total maternal deaths	22,000	(2005)






INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %) 51 (2005) Wasting prevalence (moderate and severe, %)


12 (2005)


Complementary feeding rate (6-9 months, %) Low birthweight incidence (%)


54 (2005) 20 (2005)























2 (2005)

Adolescent birth rate (births per 1,000 women) 109 (2003)

Unmet need for family planning (%) 34 (2005)


Antenatal visits for woman (4 or more visits, %) 12 (2005)

Intermittent preventive treatment for malaria (%) NA*

C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) 1, 9, 0 (2005)


Early initiation of breastfeeding (within 1 hr of birth, %) 69 (2005)


Postnatal visit for baby (within 2 days for home births, %)




2005












WATER AND SANITATION

2000





EQUITY



POLICIES

20

0

International Code of Marketing of Breastmilk Substitutes	Partial
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	Partial
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	Yes
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	No

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	30 (2007)	
General government expenditure on health as % of total government expenditure (%)	10 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	34 (2007)	
Density of health workers (per 10,000 population)	2.6 (2007)	
Official Development Assistance to child health per child (US\$)	12 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	17 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)	11 (2008)	


Ethiopia




Gabon

DEMOGRAPHICS

Total population (000)	1,448	(2008)
Total under-five population (000)	182	(2008)
Births (000)	40	(2008)
Birth registration (%)	89	(2000)
Under-five mortality rate (per 1000 live births)	77	(2008)
Infant mortality rate (per 1000 live births)	57	(2008)
Neonatal mortality rate (per 1000 live births)	28	(2008)
Total under-five deaths (000)	3	(2008)
Maternal mortality ratio (per 100,000 live births)	520	(2005)
Lifetime risk of maternal death (1 in N)	53	(2005)
Total maternal deaths	220	(2005)





INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %) 25 (2000) Wasting prevalence (moderate and severe, %)

4 (2000)


Complementary feeding rate (6-9 months, %) Low birthweight incidence (%)

62 (2000)


14 (2000)










ased on 2006 WHO reference population **CHILD HEALTH**


















Adolescent birth rate (births per 1,000 women) 144 (1998)
Unmet need for family planning (%) 28 (2000)
Antenatal visits for woman (4 or more visits, %) 63 (2000)
Intermittent preventive treatment for malaria (%) ---C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) 6,6,4 (2000)
Early initiation of breastfeeding (within 1 hr of birth, %) 71 (2000)
Postnatal visit for baby (within 2 days for home births, %) ---














WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Yes
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	Yes
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	Yes
Maternity protection in accordance with ILO Convention 183	Partial
Specific notification of maternal deaths	No

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	650 (2007)	
General government expenditure on health as % of total government expenditure (%)	14 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	36 (2007)	
Density of health workers (per 10,000 population)	53.1 (2004)	
Official Development Assistance to child health per child (US\$)	12 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	37 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)	123 (2001)	


Gabon




The Gambia

DEMOGRAPHICS

Total population (000)	1,660	(2008)
Total under-five population (000)	267	(2008)
Births (000)	61	(2008)
Birth registration (%)	55	(2005-06)
Under-five mortality rate (per 1000 live births)	106	(2008)
Infant mortality rate (per 1000 live births)	80	(2008)
Neonatal mortality rate (per 1000 live births)	34	(2008)
Total under-five deaths (000)	6	(2008)
Maternal mortality ratio (per 100,000 live births)	690	(2005)
Lifetime risk of maternal death (1 in N)	32	(2005)
Total maternal deaths	360	(2005)





INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN

Complementary feeding rate (6-9 months, %)


Stunting prevalence (moderate and severe, %) 28 (2006) Wasting prevalence (moderate and severe, %)


7 (2006)


Low birthweight incidence (%)


44 (2006)


20 (2006)



























104 (2000) Adolescent birth rate (births per 1,000 women) Unmet need for family planning (%) Antenatal visits for woman (4 or more visits, %) Intermittent preventive treatment for malaria (%) 33 (2006) C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) Early initiation of breastfeeding (within 1 hr of birth, %) 48 (2006)

















WATER AND SANITATION









POLICIES

International Code of Marketing of Breastmilk Substitutes	Yes
New ORS formula and zinc for management of diarrhoea	Partial
Community treatment of pneumonia with antibiotics	Yes
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	Partial
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	No

SYSTEMS

Financial Flows and Human Reso	ources
Per capita total expenditure on health (US\$)	90 (2007)
General government expenditure on health as % of total government expenditure (%)	12 (2007)
Out-of-pocket expenditure as % of total expenditure on health (%)	20 (2007)
Density of health workers (per 10,000 population)	6.1 (2008)
Official Development Assistance to child health per child (US\$)	20 (2007)
Official Development Assistance to maternal and neonatal health per live birth (US\$)	25 (2007)
National availability of Emergency Obstetric Care services (% of recommended minimum)	86 (2002)


The Gambia




Ghana

DEMOGRAPHICS

Total population (000)	23,351	(2008)
Total under-five population (000)	3,319	(2008)
Births (000)	757	(2008)
Birth registration (%)	71	(2008)
Under-five mortality rate (per 1000 live births)	76	(2008)
Infant mortality rate (per 1000 live births)	51	(2008)
Neonatal mortality rate (per 1000 live births)	30	(2008)
Total under-five deaths (000)	55	(2008)
Maternal mortality ratio (per 100,000 live births)	560	(2005)
Lifetime risk of maternal death (1 in N)	45	(2005)
Total maternal deaths	3,800	(2005)





INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN

NUTRITION


Stunting prevalence (moderate and severe, %) 28 (2008) Wasting prevalence (moderate and severe, %) 9 (2008)


Complementary feeding rate (6-9 months, %) Low birthweight incidence (%)


Malaria prevention


Percent children < 5 years sleeping under ITNs


75 (2008) 9 (2006)























Adolescent birth rate (births per 1,000 women) 70 (2006)

Unmet need for family planning (%) 35 (2008)

Antenatal visits for woman (4 or more visits, %) 78 (2008)


Intermittent preventive treatment for malaria (%) 44 (2008)


C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) 7, 11, 5 (2008)


Early initiation of breastfeeding (within 1 hr of birth, %) 52 (2008)


Postnatal visit for baby (within 2 days for home births, %) ---














WATER AND SANITATION







EQUITY

POLICIES

International Code of Marketing of Breastmilk Substitutes	Yes	
New ORS formula and zinc for management of diarrhoea	Yes	
Community treatment of pneumonia with antibiotics	Yes	
IMCI adapted to cover newborns 0-1 week of age	Yes	
Costed implementation plan(s) for maternal, newborn and child health available	Partial	
Midwives be authorised to administer a core set of life saving interventions	Yes	
Maternity protection in accordance with ILO Convention 183	Partial	
Specific notification of maternal deaths	Yes	

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	113	(2007)
General government expenditure on health as % of total government expenditure (%)	11	(2007)
Out-of-pocket expenditure as % of total expenditure on health (%)	38	(2007)
Density of health workers (per 10,000 population)	10.8	(2008)
Official Development Assistance to child health per child (US\$)	21	(2007)
Official Development Assistance to maternal an neonatal health per live birth (US\$)		(2007)
National availability of Emergency Obstetric Ca services (% of recommended minimum)		(2004-2005)


Ghana




Guatemala

DEMOGRAPHICS

Total population (000)	13,686	(2008)
Total under-five population (000)	2,118	(2008)
Births (000)	453	(2008)
Birth registration (%)		
Under-five mortality rate (per 1000 live births)	35	(2008)
Infant mortality rate (per 1000 live births)	29	(2008)
Neonatal mortality rate (per 1000 live births)	11	(2008)
Total under-five deaths (000)	15	(2008)
Maternal mortality ratio (per 100,000 live births)	290	(2005)
Lifetime risk of maternal death (1 in N)	71	(2005)
Total maternal deaths	1,300	(2005)







INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %)

54 (2002) 2 (2002) Complementary feeding rate (6-9 months, %) Low birthweight incidence (%) 67 (2002) 12 (2002)






















Adolescent birth rate (births per 1,000 women)

92 (2006)

Unmet need for family planning (%)

Antenatal visits for woman (4 or more visits, %)

Intermittent preventive treatment for malaria (%)

NA*


C-section rate (total, urban, rural; %)

(Minimum target is 5% and maximum target is 15%) 11,19,8 (2002)


Early initiation of breastfeeding (within 1 hr of birth, %) 60 (2002)


Postnatal visit for baby (within 2 days for home births, %)

















WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Yes
New ORS formula and zinc for management of diarrhoea	Partial
Community treatment of pneumonia with antibiotics	Yes
IMCI adapted to cover newborns 0-1 week of age	Partial
Costed implementation plan(s) for maternal, newborn and child health available	Partial
Midwives be authorised to administer a core set of life saving interventions	Partial
Maternity protection in accordance with ILO Convention 183	Partial
Specific notification of maternal deaths	Partial

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	334 (2007)	
General government expenditure on health as % of total government expenditure (%)	14 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	66 (2007)	
Density of health workers (per 10,000 population)	49.5 (1999)	
Official Development Assistance to child health per child (US\$)	7 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	15 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)		


Guatemala




Guinea

DEMOGRAPHICS

Total population (000)	9,833	(2008)
Total under-five population (000)	1,635	(2008)
Births (000)	392	(2008)
Birth registration (%)	43	(2005)
Under-five mortality rate (per 1000 live births)	146	(2008)
Infant mortality rate (per 1000 live births)	90	(2008)
Neonatal mortality rate (per 1000 live births)	43	(2008)
Total under-five deaths (000)	54	(2008)
Maternal mortality ratio (per 100,000 live births)	910	(2005)
Lifetime risk of maternal death (1 in N)	19	(2005)
Total maternal deaths	3,500	(2005)






INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %) Wasting prevalence (moderate and severe, %)


40 (2008) 8 (2008) Complementary feeding rate (6-9 months, %) Low birthweight incidence (%)


32 (2008) 12 (2005)























Adolescent birth rate (births per 1,000 women) 153 (2003)

Unmet need for family planning (%) 21 (2005)

Antenatal visits for woman (4 or more visits, %) 50 (2007)


Intermittent preventive treatment for malaria (%) 3 (2005)


C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) 2, 5, 2 (2007)

Early initiation of breastfeeding (within 1 hr of birth, %) 35 (2008)


Postnatal visit for baby (within 2 days for home births, %) ---














WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Partial
New ORS formula and zinc for management of diarrhoea	No
Community treatment of pneumonia with antibiotics	No
IMCI adapted to cover newborns 0-1 week of age	No
Costed implementation plan(s) for maternal, newborn and child health available	No
Midwives be authorised to administer a core set of life saving interventions	Partial
Maternity protection in accordance with ILO Convention 183	Partial
Specific notification of maternal deaths	Partial

SYSTEMS

Financial Flows and Human Reso	ources
Per capita total expenditure on health (US\$)	62 (2007)
General government expenditure on health as % of total government expenditure (%)	5 (2007)
Out-of-pocket expenditure as % of total expenditure on health (%)	89 (2007)
Density of health workers (per 10,000 population)	1.4 (2005)
Official Development Assistance to child health per child (US\$)	4 (2007)
Official Development Assistance to maternal and neonatal health per live birth (US\$)	8 (2007)
National availability of Emergency Obstetric Care services (% of recommended minimum)	25 (2002)


Guinea




Guinea-Bissau

DEMOGRAPHICS

Total population (000)	1,575	(2008)
Total under-five population (000)	265	(2008)
Births (000)	65	(2008)
Birth registration (%)	39	(2006)
Under-five mortality rate (per 1000 live births)	195	(2008)
Infant mortality rate (per 1000 live births)	117	(2008)
Neonatal mortality rate (per 1000 live births)	45	(2008)
Total under-five deaths (000)	12	(2008)
Maternal mortality ratio (per 100,000 live births)	1,100	(2005)
Lifetime risk of maternal death (1 in N)	13	(2005)
Total maternal deaths	890	(2005)






INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %)


47 (2006) Wasting prevalence (moderate and severe, %) 8 (2006)


Complementary feeding rate (6-9 months, %) Low birthweight incidence (%)


35 (2006) 24 (2006)



























Postnatal visit for baby (within 2 days for home births, %)














WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Partial
New ORS formula and zinc for management of diarrhoea	Partial
Community treatment of pneumonia with antibiotics	No
IMCI adapted to cover newborns 0-1 week of age	No
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	No
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	Yes

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	33 (2007)	
General government expenditure on health as % of total government expenditure (%)	4 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	41 (2007)	
Density of health workers (per 10,000 population)	6.0 (2008)	
Official Development Assistance to child health per child (US\$)	12 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	20 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)	127 (2002)	


Guinea-Bissau




Haiti

DEMOGRAPHICS

Total population (000)	9,876	(2008)
Total under-five population (000)	1,252	(2008)
Births (000)	273	(2008)
Birth registration (%)	81	(2005)
Under-five mortality rate (per 1000 live births)	72	(2008)
Infant mortality rate (per 1000 live births)	54	(2008)
Neonatal mortality rate (per 1000 live births)	25	(2008)
Total under-five deaths (000)	19	(2008)
Maternal mortality ratio (per 100,000 live births)	670	(2005)
Lifetime risk of maternal death (1 in N)	44	(2005)
Total maternal deaths	1,700	(2005)








INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %) 29 (2005-2006) Wasting prevalence (moderate and severe, %) 10 (2005-2006)


Complementary feeding rate (6-9 months, %) 87 (2005-2006) Low birthweight incidence (%) 25 (2005-2006)























4

Adolescent birth rate (births per 1,000 women) 69 (2003)

Unmet need for family planning (%) 38 (2005-06)


Antenatal visits for woman (4 or more visits, %) 54 (2005-06)


Intermittent preventive treatment for malaria (%) --
C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) 3, 6, 1 (2005)

Early initiation of breastfeeding (within 1 hr of birth, %) 44 (2005-06)


Postnatal visit for baby (within 2 days for home births, %)














WATER AND SANITATION









POLICIES

International Code of Marketing of Breastmilk Substitutes	Partial
New ORS formula and zinc for management of diarrhoea	No
Community treatment of pneumonia with antibiotics	No
IMCI adapted to cover newborns 0-1 week of age	No
Costed implementation plan(s) for maternal, newborn and child health available	No
Midwives be authorised to administer a core set of life saving interventions	No
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	No

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	58 (2007)	
General government expenditure on health as % of total government expenditure (%)	9 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	44 (2007)	
Density of health workers (per 10,000 population)	3.6 (1998)	
Official Development Assistance to child health per child (US\$)	16 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	54 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)		


Haiti




India

DEMOGRAPHICS

Total population (000) 1,181,412 (2008) Total under-five population (000) 126,642 (2008) 26,913 (2008) Births (000) Birth registration (%) 41 (2006) Under-five mortality rate (per 1000 live births) 69 (2008) Infant mortality rate (per 1000 live births) 52 (2008) Neonatal mortality rate (per 1000 live births) 37 (2008) Total under-five deaths (000) 1,830 (2008) 450 (2005) Maternal mortality ratio (per 100,000 live births) Lifetime risk of maternal death (1 in N) 70 (2005) 117,000 (2005) Total maternal deaths







INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN

Stunting prevalence (moderate and severe, %) 48 (2005-2006) Wasting prevalence (moderate and severe, %) 20 (2005-2006)


Complementary feeding rate (6-9 months, %) 57 (2005-2006) Low birthweight incidence (%) 28 (2005-2006)











*Based on 2006 WHO reference population














Adolescent birth rate (births per 1,000 women) 45 (2006)
Unmet need for family planning (%) 13 (2005-06)
Antenatal visits for woman (4 or more visits, %) 37 (2005-06)
Intermittent preventive treatment for malaria (%) NA*


Intermittent preventive treatment for maiaria (%)

C-section rate (total, urban, **rural**; %) (Minimum target is 5% and maximum target is 15%) 9,17,**6** (2005-06)


Early initiation of breastfeeding (within 1 hr of birth, %) 25 (2005-06)


Postnatal visit for baby (within 2 days for home births, %) ---


^{*} Not applicable














WATER AND SANITATION





EQUITY



POLICIES

Yes
Yes
Yes
Yes
Yes
Partial
No
Yes

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	109 (2007)	
General government expenditure on health as % of total government expenditure (%)	4 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	66 (2007)	
Density of health workers (per 10,000 population)	18.5 (2005)	
Official Development Assistance to child health per child (US\$)	1 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	5 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)		


India




Indonesia

DEMOGRAPHICS

Total population (000)	227,345	(2008)
Total under-five population (000)	20,891	(2008)
Births (000)	4,220	(2008)
Birth registration (%)	55	(2002)
Under-five mortality rate (per 1000 live births) 41	(2008)
Infant mortality rate (per 1000 live births)	31	(2008)
Neonatal mortality rate (per 1000 live births)	19	(2008)
Total under-five deaths (000)	173	(2008)
Maternal mortality ratio (per 100,000 live births	420	(2005)
Lifetime risk of maternal death (1 in N)	97	(2005)
Total maternal deaths	19,000	(2005)






INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN

Stunting prevalence (moderate and severe, %) 37 (2007) Wasting prevalence (moderate and severe, %)


14 (2007)

Complementary feeding rate (6-9 months, %) 75 (2007)


Low birthweight incidence (%) 9 (2007)











Based on 2006 WHO reference population














Adolescent birth rate (births per 1,000 women) 52 (2005)

Unmet need for family planning (%) 9 (2007)

Antenatal visits for woman (4 or more visits, %) 82 (2007)


Intermittent preventive treatment for malaria (%) NA*

C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) 7,11,4 (2007)


Early initiation of breastfeeding (within 1 hr of birth, %) 39 (2007)


Postnatal visit for baby (within 2 days for home births, %) ---

















WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Partial
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	Partial
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	No
Midwives be authorised to administer a core set of life saving interventions	Partial
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	No

SYSTEMS


Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	81 (2007)	
General government expenditure on health as % of total government expenditure (%)	6 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	30 (2007)	
Density of health workers (per 10,000 population)	9.5 (2003)	
Official Development Assistance to child health per child (US\$)	2 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	6 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)		


Indonesia

Iraq

DEMOGRAPHICS

Total population (000)	30,096	(2008)
Total under-five population (000)	4,450	(2008)
Births (000)	944	(2008)
Birth registration (%)	95	(2006)
Under-five mortality rate (per 1000 live births)	44	(2008)
Infant mortality rate (per 1000 live births)	36	(2008)
Neonatal mortality rate (per 1000 live births)	25	(2008)
Total under-five deaths (000)	41	(2008)
Maternal mortality ratio (per 100,000 live births)	300	(2005)
Lifetime risk of maternal death (1 in N)	72	(2005)
Total maternal deaths	2,900	(2005)






INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %) 26 (2006) Wasting prevalence (moderate and severe, %)

6 (2006)

Complementary feeding rate (6-9 months, %) Low birthweight incidence (%)


51 (2006) 15 (2006)











Based on 2006 WHO reference population














Adolescent birth rate (births per 1,000 women) 68 (2005)


Unmet need for family planning (%) --
Antenatal visits for woman (4 or more visits, %) --
Intermittent preventive treatment for malaria (%) NA*


C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) 21, 24, 15 (2006)


Early initiation of breastfeeding (within 1 hr of birth, %) 31 (2006)


Postnatal visit for baby (within 2 days for home births, %) ---


^{*} Not applicable













WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Partial
New ORS formula and zinc for management of diarrhoea	Partial
Community treatment of pneumonia with antibiotics	No
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	No
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	Yes

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	78 (2007)	
General government expenditure on health as % of total government expenditure (%)	3 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	25 (2007)	
Density of health workers (per 10,000 population)	15.8 (2007)	
Official Development Assistance to child health per child (US\$)	12 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	30 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)		


Iraq




Kenya

DEMOGRAPHICS

Total population (000)	38,765	(2008)
Total under-five population (000)	6,540	(2008)
Births (000)	1,506	(2008)
Birth registration (%)	48	(2003)
Under-five mortality rate (per 1000 live births)	128	(2008)
Infant mortality rate (per 1000 live births)	81	(2008)
Neonatal mortality rate (per 1000 live births)	33	(2008)
Total under-five deaths (000)	189	(2008)
Maternal mortality ratio (per 100,000 live births)	560	(2005)
Lifetime risk of maternal death (1 in N)	39	(2005)
Total maternal deaths	7,700	(2005)







INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %) 35 (2008-2009)


Wasting prevalence (moderate and severe, %) 7 (2008-2009)





Complementary feeding rate (6-9 months, %) 83 (2008-2009) Low birthweight incidence (%) (2003)






















103 (2008)Adolescent birth rate (births per 1,000 women) Unmet need for family planning (%) 25 (2003)Antenatal visits for woman (4 or more visits, %) 52 (2003) Intermittent preventive treatment for malaria (%) 15 (2008-09) C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) 4, 9, 3 (2003)Early initiation of breastfeeding (within 1 hr of birth, %) 52 (2003)















WATER AND SANITATION


1998

2000


2003 2008-09

1993





EQUITY



POLICIES

100

80

60

40

20

1989

International Code of Marketing of Breastmilk Substitutes	Partial
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	No
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	Yes
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	Yes

SYSTEMS


Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	72 (2007)	
General government expenditure on health as % of total government expenditure (%)	8 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	45 (2007)	
Density of health workers (per 10,000 population)	13.2 (2002)	
Official Development Assistance to child health per child (US\$)	10 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	14 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)		


Kenya

Korea, Democratic People's Republic of

DEMOGRAPHICS

Total population (000)	23,819	(2008)
Total under-five population (000)	1,575	(2008)
Births (000)	327	(2008)
Birth registration (%)	99	(2000)
Under-five mortality rate (per 1000 live births)	55	(2008)
Infant mortality rate (per 1000 live births)	42	(2008)
Neonatal mortality rate (per 1000 live births)	29	(2008)
Total under-five deaths (000)	18	(2008)
Maternal mortality ratio (per 100,000 live births)	370	(2005)
Lifetime risk of maternal death (1 in N)	140	(2005)
Total maternal deaths	1,300	(2005)






INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %) 45 (2004) Wasting prevalence (moderate and severe, %)


9 (2004)

Complementary feeding rate (6-9 months, %) Low birthweight incidence (%)


31 (2004) 7 (2002)










ased on 2006 WHO reference population














Adolescent birth rate (births per 1,000 women) 0 (1994)


Unmet need for family planning (%) --
Antenatal visits for woman (4 or more visits, %) --
Intermittent preventive treatment for malaria (%) NA*


C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) ---, ---, --
Early initiation of breastfeeding (within 1 hr of birth, %) --
Postnatal visit for baby (within 2 days for home births, %) ---


^{*} Not applicable













WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	No
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	Yes
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Partial
Midwives be authorised to administer a core set of life saving interventions	Partial
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	Partial

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)		
General government expenditure on health as % of total government expenditure (%)		
Out-of-pocket expenditure as % of total expenditure on health (%)		
Density of health workers (per 10,000 population)	74.1 (2003)	
Official Development Assistance to child health per child (US\$)	3 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	9 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)		


Korea, Democratic People's Republic of




Lao People's Democratic Republic

DEMOGRAPHICS

Total population (000)	6,205	(2008)
Total under-five population (000)	776	(2008)
Births (000)	170	(2008)
Birth registration (%)	72	(2006)
Under-five mortality rate (per 1000 live births)	61	(2008)
Infant mortality rate (per 1000 live births)	48	(2008)
Neonatal mortality rate (per 1000 live births)	20	(2008)
Total under-five deaths (000)	10	(2008)
Maternal mortality ratio (per 100,000 live births)	660	(2005)
Lifetime risk of maternal death (1 in N)	33	(2005)
Total maternal deaths	1,300	(2005)






INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN

Stunting prevalence (moderate and severe, %) 48 (2006) Wasting prevalence (moderate and severe, %)


7 (2006)


Complementary feeding rate (6-9 months, %) Low birthweight incidence (%)


70 (2006) 11 (2006)























Adolescent birth rate (births per 1,000 women) 110 (2005)


Unmet need for family planning (%) 40 (2000)


Antenatal visits for woman (4 or more visits, %) --
Intermittent preventive treatment for malaria (%) 1 (2006)

C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) ---, ---, --
Early initiation of breastfeeding (within 1 hr of birth, %) 30 (2006)


Postnatal visit for baby (within 2 days for home births, %) ---













WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes Partial New ORS formula and zinc for management of diarrhoea Partial Community treatment of pneumonia with antibiotics IMCI adapted to cover newborns 0-1 week of age No Costed implementation plan(s) for maternal, newborn and child health available Yes Midwives be authorised to administer a core set of life saving interventions Partial Maternity protection in accordance with ILO Convention 183 No Specific notification of maternal deaths Partial		
diarrhoea Partial Community treatment of pneumonia with antibiotics Partial IMCl adapted to cover newborns 0-1 week of age No Costed implementation plan(s) for maternal, newborn and child health available Yes Midwives be authorised to administer a core set of life saving interventions Partial Maternity protection in accordance with ILO Convention 183 No	Ŭ	Partial
IMCI adapted to cover newborns 0-1 week of age Costed implementation plan(s) for maternal, newborn and child health available Yes Midwives be authorised to administer a core set of life saving interventions Partial Maternity protection in accordance with ILO Convention 183 No		Partial
Costed implementation plan(s) for maternal, newborn and child health available Midwives be authorised to administer a core set of life saving interventions Partial Maternity protection in accordance with ILO Convention 183 No	Community treatment of pneumonia with antibiotics	Partial
newborn and child health available Yes Midwives be authorised to administer a core set of life saving interventions Partial Maternity protection in accordance with ILO Convention 183 No	IMCI adapted to cover newborns 0-1 week of age	No
life saving interventions Partial Maternity protection in accordance with ILO Convention 183 No		Yes
Convention 183 No		Partial
Specific notification of maternal deaths Partial		No
	Specific notification of maternal deaths	Partial

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)		
Per capita total experiolitire of Friedli (105\$)	84 (2007)	
General government expenditure on health as % of total government expenditure (%)	4 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	62 (2007)	
Density of health workers (per 10,000 population)	13.2 (2004)	
Official Development Assistance to child health per child (US\$)	7 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	25 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)		


Lao People's Democratic Republic




Lesotho

DEMOGRAPHICS

Total population (000)	2,049	(2008)
Total under-five population (000)	272	(2008)
Births (000)	59	(2008)
Birth registration (%)	26	(2004)
Under-five mortality rate (per 1000 live births)	79	(2008)
Infant mortality rate (per 1000 live births)	63	(2008)
Neonatal mortality rate (per 1000 live births)	37	(2008)
Total under-five deaths (000)	5	(2008)
Maternal mortality ratio (per 100,000 live births)	960	(2005)
Lifetime risk of maternal death (1 in N)	45	(2005)
Total maternal deaths	480	(2005)





INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Complementary feeding rate (6-9 months, %)


Stunting prevalence (moderate and severe, %) 42 (2007) Wasting prevalence (moderate and severe, %)


2 (2007)

Low birthweight incidence (%)


79 (2004) 13 (2004)











Based on 2006 WHO reference population


















Adolescent birth rate (births per 1,000 women)	98	(2003)
Unmet need for family planning (%)	31	(2004)
Antenatal visits for woman (4 or more visits, %)	70	(2004)
Intermittent preventive treatment for malaria (%)		
C-section rate (total, urban, rural ; %) (Minimum target is 5% and maximum target is 15%)	5, 8, 5	(2004)
Early initiation of breastfeeding (within 1 hr of birth, %)	63	(2004)
Postnatal visit for baby (within 2 days for home births, %))	














WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	No
New ORS formula and zinc for management of diarrhoea	Partial
Community treatment of pneumonia with antibiotics	No
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Partial
Midwives be authorised to administer a core set of life saving interventions	Partial
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	Yes

SYSTEMS

Financial Flows and Human Resources		
92	(2007)	
8	(2007)	
29	(2007)	
6.7	(2003)	
10	(2007)	
21	(2007)	
29	(2004)	
	92 8 29 6.7 10 21	


Lesotho




Liberia

DEMOGRAPHICS

Total population (000)	3,793	(2008)
Total under-five population (000)	619	(2008)
Births (000)	145	(2008)
Birth registration (%)	4	(2007)
Under-five mortality rate (per 1000 live births)	145	(2008)
Infant mortality rate (per 1000 live births)	100	(2008)
Neonatal mortality rate (per 1000 live births)	44	(2008)
Total under-five deaths (000)	20	(2008)
Maternal mortality ratio (per 100,000 live births)	1,200	(2005)
Lifetime risk of maternal death (1 in N)	12	(2005)
Total maternal deaths	2,100	(2005)





INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN

NUTRITION

100

80

60

40

20

0

Underweight prevalence

Percent children < 5 years underweight for age*


23

Other NS


19

2007


Stunting prevalence (moderate and severe, %) 39 (2007) Wasting prevalence (moderate and severe, %) 8 (2007) Complementary feeding rate (6-9 months, %)
Low birthweight incidence (%)





62 (2007)





*Based on 2006 WHO reference population


















Adolescent birth rate (births per 1,000 women)	177	(2007)
Unmet need for family planning (%)	36	(2007)
Antenatal visits for woman (4 or more visits, %)	66	(2007)
Intermittent preventive treatment for malaria (%)	45	(2009)
C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%)	4, 5, 3	(2007)
Early initiation of breastfeeding (within 1 hr of birth, %)	67	(2007)
Postnatal visit for haby (within 2 days for home births 9	۵)	














WATER AND SANITATION









POLICIES

No
Yes
Partial
Yes
Yes
Yes
No
Partial

SYSTEMS

Financial Flows and Human Reso	ources
Per capita total expenditure on health (US\$)	39 (2007)
General government expenditure on health as % of total government expenditure (%)	17 (2007)
Out-of-pocket expenditure as % of total expenditure on health (%)	39 (2007)
Density of health workers (per 10,000 population)	2.8 (2008)
Official Development Assistance to child health per child (US\$)	22 (2007)
Official Development Assistance to maternal and neonatal health per live birth (US\$)	67 (2007)
National availability of Emergency Obstetric Care services (% of recommended minimum)	


Liberia




Madagascar

DEMOGRAPHICS

Total population (000)	19,111	(2008)
Total under-five population (000)	3,060	(2008)
Births (000)	687	(2008)
Birth registration (%)	75	(2003-04)
Under-five mortality rate (per 1000 live births)	106	(2008)
Infant mortality rate (per 1000 live births)	68	(2008)
Neonatal mortality rate (per 1000 live births)	35	(2008)
Total under-five deaths (000)	71	(2008)
Maternal mortality ratio (per 100,000 live births)	510	(2005)
Lifetime risk of maternal death (1 in N)	38	(2005)
Total maternal deaths	3,600	(2005)








INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %) 53 (2003-2004) Wasting prevalence (moderate and severe, %) 15 (2003-2004)


Complementary feeding rate (6-9 months, %) 78 (2003-2004) Low birthweight incidence (%) 17 (2003-2004)























Adolescent birth rate (births per 1,000 women)

148 (2008)

Unmet need for family planning (%)

24 (2003-04)

Antenatal visits for woman (4 or more visits, %)


Intermittent preventive treatment for malaria (%)

6 (2008-09)


C-section rate (total, urban, **rural**; %)
(Minimum target is 5% and maximum target is 15%)
1,2,1 (2003-04)


Early initiation of breastfeeding (within 1 hr of birth, %) 62 (2003-04)


Postnatal visit for baby (within 2 days for home births, %) ---














WATER AND SANITATION





EQUITY



POLICIES

	International Code of Marketing of Breastmilk Substitutes	Yes
	New ORS formula and zinc for management of diarrhoea	Yes
	Community treatment of pneumonia with antibiotics	Yes
	IMCI adapted to cover newborns 0-1 week of age	Yes
	Costed implementation plan(s) for maternal, newborn and child health available	Yes
	Midwives be authorised to administer a core set of life saving interventions	Yes
	Maternity protection in accordance with ILO Convention 183	No
	Specific notification of maternal deaths	Partial
ı		

SYSTEMS

Financial Flows and Human Re	sources
Per capita total expenditure on health (US\$)	41 (2007)
General government expenditure on health as % of total government expenditure (%)	15 (2007)
Out-of-pocket expenditure as % of total expenditure on health (%)	23 (2007)
Density of health workers (per 10,000 population)	4.8 (2007)
Official Development Assistance to child health per child (US\$)	10 (2007)
Official Development Assistance to maternal and neonatal health per live birth (US\$)	d 17 (2007)
National availability of Emergency Obstetric Card services (% of recommended minimum)	e 30 (2002-2003)


Madagascar




Malawi

DEMOGRAPHICS

Total population (000)	14,846	(2008)
Total under-five population (000)	2,591	(2008)
Births (000)	599	(2008)
Birth registration (%)		
Under-five mortality rate (per 1000 live births)	100	(2008)
Infant mortality rate (per 1000 live births)	65	(2008)
Neonatal mortality rate (per 1000 live births)	29	(2008)
Total under-five deaths (000)	56	(2008)
Maternal mortality ratio (per 100,000 live births)	1,100	(2005)
Lifetime risk of maternal death (1 in N)	18	(2005)
Total maternal deaths	6,000	(2005)





INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN

Stunting prevalence (moderate and severe, %) 53 (2006) Wasting prevalence (moderate and severe, %)


4 (2006)


Low birthweight incidence (%)


Complementary feeding rate (6-9 months, %)

89 (2006)


14 (2006)











Based on 2006 WHO reference population














Adolescent birth rate (births per 1,000 women) 178 (2005)

Unmet need for family planning (%) 28 (2004)

Antenatal visits for woman (4 or more visits, %) 57 (2004)


Intermittent preventive treatment for malaria (%) 45 (2006)


C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) 3, 4, 3 (2004)


Early initiation of breastfeeding (within 1 hr of birth, %) 58 (2006)


Postnatal visit for baby (within 2 days for home births, %) 3 (2006)














WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Yes
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	Yes
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	Yes
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	Partial

SYSTEMS

Financial Flows and Human Reso	ources
Per capita total expenditure on health (US\$)	50 (2007)
General government expenditure on health as % of total government expenditure (%)	12 (2007)
Out-of-pocket expenditure as % of total expenditure on health (%)	11 (2007)
Density of health workers (per 10,000 population)	3.0 (2008)
Official Development Assistance to child health per child (US\$)	19 (2007)
Official Development Assistance to maternal and neonatal health per live birth (US\$)	35 (2007)
National availability of Emergency Obstetric Care services (% of recommended minimum)	37 (2004)


Malawi




Mali

DEMOGRAPHICS

Total population (000)	12,706	(2008)
Total under-five population (000)	2,207	(2008)
Births (000)	542	(2008)
Birth registration (%)	53	(2006)
Under-five mortality rate (per 1000 live births)	194	(2008)
Infant mortality rate (per 1000 live births)	103	(2008)
Neonatal mortality rate (per 1000 live births)	52	(2008)
Total under-five deaths (000)	100	(2008)
Maternal mortality ratio (per 100,000 live births)	970	(2005)
Lifetime risk of maternal death (1 in N)	15	(2005)
Total maternal deaths	6,400	(2005)





INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN

Stunting prevalence (moderate and severe, %) 38 (2006) Wasting prevalence (moderate and severe, %)


15 (2006)


Low birthweight incidence (%)


Complementary feeding rate (6-9 months, %)

30 (2006)


19 (2006)











Based on 2006 WHO reference population


















Adolescent birth rate (births per 1,000 women)	190	(2004)	
Unmet need for family planning (%)	31	(2006)	
,, ,,	01	,	
Antenatal visits for woman (4 or more visits, %)	35	(2006)	
Intermittent preventive treatment for malaria (%)	4	(2006)	
C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%)	2, 4, 1	(2006)	
Early initiation of breastfeeding (within 1 hr of birth, %)	46	(2006)	
Postnatal visit for baby (within 2 days for home births %	3		














WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Partial
New ORS formula and zinc for management of diarrhoea	Partial
Community treatment of pneumonia with antibiotics	Yes
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	Partial
Maternity protection in accordance with ILO Convention 183	Yes
Specific notification of maternal deaths	Yes

SYSTEMS

Financial Flows and Human Resources Per capita total expenditure on health (US\$) 67 (2007) General government expenditure on health as % of total government expenditure (%) 12 (2007) Out-of-pocket expenditure as % of total expenditure on health (%) 48 (2007) Density of health workers (per 10,000 population) 2.7 (2007) Official Development Assistance to child health per child (US\$) 13 (2007) Official Development Assistance to maternal and neonatal health per live birth (US\$) 23 (2007) National availability of Emergency Obstetric Care services (% of recommended minimum) 38 (2003)		
General government expenditure on health as % of total government expenditure (%) 12 (2007) Out-of-pocket expenditure as % of total expenditure on health (%) 48 (2007) Density of health workers (per 10,000 population) 2.7 (2007) Official Development Assistance to child health per child (US\$) 13 (2007) Official Development Assistance to maternal and neonatal health per live birth (US\$) 23 (2007) National availability of Emergency Obstetric Care	Financial Flows and Human Reso	ources
% of total government expenditure (%) Out-of-pocket expenditure as % of total expenditure on health (%) Density of health workers (per 10,000 population) Official Development Assistance to child health per child (US\$) Official Development Assistance to maternal and neonatal health per live birth (US\$) National availability of Emergency Obstetric Care	Per capita total expenditure on health (US\$)	67 (2007)
expenditure on health (%) Density of health workers (per 10,000 population) Official Development Assistance to child health per child (US\$) Official Development Assistance to maternal and neonatal health per live birth (US\$) National availability of Emergency Obstetric Care		12 (2007)
Official Development Assistance to child health per child (US\$) 13 (2007) Official Development Assistance to maternal and neonatal health per live birth (US\$) 23 (2007) National availability of Emergency Obstetric Care		48 (2007)
per child (US\$) 13 (2007) Official Development Assistance to maternal and neonatal health per live birth (US\$) 23 (2007) National availability of Emergency Obstetric Care	Density of health workers (per 10,000 population)	2.7 (2007)
neonatal health per live birth (US\$) 23 (2007) National availability of Emergency Obstetric Care		13 (2007)
, , ,		23 (2007)
	, , ,	38 (2003)


Mali




Mauritania

DEMOGRAPHICS

Total population (000)	3,215	(2008)
Total under-five population (000)	475	(2008)
Births (000)	108	(2008)
Birth registration (%)	56	(2007)
Under-five mortality rate (per 1000 live births)	118	(2008)
Infant mortality rate (per 1000 live births)	75	(2008)
Neonatal mortality rate (per 1000 live births)	45	(2008)
Total under-five deaths (000)	12	(2008)
Maternal mortality ratio (per 100,000 live births)	820	(2005)
Lifetime risk of maternal death (1 in N)	22	(2005)
Total maternal deaths	1,000	(2005)








INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN

Stunting prevalence (moderate and severe, %) 32 (2008) Wasting prevalence (moderate and severe, %) 12 (2008)


Complementary feeding rate (6-9 months, %) 72 (2008) Low birthweight incidence (%) 34 (2007)











CHILD HEALTH













10

2006

2007

2008

2005

2000-2001 2007


Adolescent birth rate (births per 1,000 women) 88 (2002)
Unmet need for family planning (%) 32 (2000-01)
Antenatal visits for woman (4 or more visits, %) 16 (2000-01)
Intermittent preventive treatment for malaria (%) ---

C-section rate (total, urban, **rural**; %)


 $(Minimum\ target\ is\ 5\%\ and\ maximum\ target\ is\ 15\%) \\ \qquad 3,6, \textbf{1} \quad (2000-01)$


Early initiation of breastfeeding (within 1 hr of birth, %) 60 (2008)


Postnatal visit for baby (within 2 days for home births, %) ---













WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	No
New ORS formula and zinc for management of diarrhoea	No
Community treatment of pneumonia with antibiotics	No
IMCI adapted to cover newborns 0-1 week of age	No
Costed implementation plan(s) for maternal, newborn and child health available	No
Midwives be authorised to administer a core set of life saving interventions	Yes
Maternity protection in accordance with ILO Convention 183	Partial
Specific notification of maternal deaths	Partial

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	47 (2007)	
General government expenditure on health as % of total government expenditure (%)	5 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	35 (2007)	
Density of health workers (per 10,000 population)	8.0 (2009)	
Official Development Assistance to child health per child (US\$)	6 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	27 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)	31 (2000)	


Mauritania




Mexico

DEMOGRAPHICS

Total population (000)	108,555	(2008)
Total under-five population (000)	10,281	(2008)
Births (000)	2,049	(2008)
Birth registration (%)		
Under-five mortality rate (per 1000 live births)	17	(2008)
Infant mortality rate (per 1000 live births)	15	(2008)
Neonatal mortality rate (per 1000 live births)	7	(2008)
Total under-five deaths (000)	36	(2008)
Maternal mortality ratio (per 100,000 live births	60	(2005)
Lifetime risk of maternal death (1 in N)	670	(2005)
Total maternal deaths	1,300	(2005)







INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


NUTRITION

Stunting prevalence (moderate and severe, %) ---Wasting prevalence (moderate and severe, %) ----

Complementary feeding rate (6-9 months, %) 36 (1987) Low birthweight incidence (%) 8 (2006)










*Based on 2006 WHO reference population

CHILD HEALTH













Pneumonia treatment Percent children < 5 years with suspected pneumonia taken to appropriate health provider Percent children < 5 years with suspected pneumonia receiving antibiotics

No data

Adolescent birth rate (births per 1,000 women) 90 (2008)

Unmet need for family planning (%) 12 (2006)


Antenatal visits for woman (4 or more visits, %) ---
Intermittent preventive treatment for malaria (%) NA*

C-section rate (total, urban, rural; %)
(Minimum target is 5% and maximum target is 15%) 38, ---, --- (2006)


Early initiation of breastfeeding (within 1 hr of birth, %) ----


Postnatal visit for baby (within 2 days for home births, %)


^{*} Not applicable












WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Partial
New ORS formula and zinc for management of diarrhoea	Partial
Community treatment of pneumonia with antibiotics	Yes
IMCI adapted to cover newborns 0-1 week of age	No
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	No
Maternity protection in accordance with ILO Convention 183	Partial
Specific notification of maternal deaths	Yes

SYSTEMS

Financial Flows and Human Resources			
Per capita total expenditure on health (US\$)	819 (2007)		
General government expenditure on health as % of total government expenditure (%)	16 (2007)		
Out-of-pocket expenditure as % of total expenditure on health (%)	51 (2007)		
Density of health workers (per 10,000 population)	68.7 (2004)		
Official Development Assistance to child health per child (US\$)	0 (2007)		
Official Development Assistance to maternal and neonatal health per live birth (US\$)	1 (2007)		
National availability of Emergency Obstetric Care services (% of recommended minimum)			


Mexico




Morocco

DEMOGRAPHICS

Total population (000)	31,606	(2008)
Total under-five population (000)	3,041	(2008)
Births (000)	646	(2006)
Birth registration (%)	85	(2000)
Under-five mortality rate (per 1000 live births)	36	(2008)
Infant mortality rate (per 1000 live births)	32	(2008)
Neonatal mortality rate (per 1000 live births)	23	(2008)
Total under-five deaths (000)	24	(2008)
Maternal mortality ratio (per 100,000 live births)	240	(2005)
Lifetime risk of maternal death (1 in N)	150	(2005)
Total maternal deaths	1,700	(2005)








INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN

Stunting prevalence (moderate and severe, %) 23 (2003-2004) Wasting prevalence (moderate and severe, %) 10 (2003-2004)


Complementary feeding rate (6-9 months, %) 66 (2003-2004) Low birthweight incidence (%) 15 (2003-2004)











Based on 2006 WHO reference population















Adolescent birth rate (births per 1,000 women) 18 (2005)
Unmet need for family planning (%) 10 (2003-04)
Antenatal visits for woman (4 or more visits, %) 31 (2003-04)
Intermittent preventive treatment for malaria (%) NA*

C-section rate (total, urban, **rural**; %) (Minimum target is 5% and maximum target is 15%) 5,9,**2** (2003-04)


Early initiation of breastfeeding (within 1 hr of birth, %) 52 (2003-04)


Postnatal visit for baby (within 2 days for home births, %) ---


* Not applicable














WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	No
New ORS formula and zinc for management of diarrhoea	No
Community treatment of pneumonia with antibiotics	No
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Partial
Midwives be authorised to administer a core set of life saving interventions	
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	Yes

SYSTEMS

Financial Flows and Human Resources			
Per capita total expenditure on health (US\$)	202 (2007)		
General government expenditure on health as % of total government expenditure (%)	6 (2007)		
Out-of-pocket expenditure as % of total expenditure on health (%)	57 (2007)		
Density of health workers (per 10,000 population)	13.4 (2007)		
Official Development Assistance to child health per child (US\$)	3 (2007)		
Official Development Assistance to maternal and neonatal health per live birth (US\$)	16 (2007)		
National availability of Emergency Obstetric Care services (% of recommended minimum)	69 (2000)		


Morocco




Mozambique

DEMOGRAPHICS

Total population (000)	22,383	(2008)
Total under-five population (000)	3,820	(2008)
Births (000)	876	(2008)
Birth registration (%)	31	(2008)
Under-five mortality rate (per 1000 live births)	130	(2008)
Infant mortality rate (per 1000 live births)	90	(2008)
Neonatal mortality rate (per 1000 live births)	43	(2008)
Total under-five deaths (000)	110	(2008)
Maternal mortality ratio (per 100,000 live births)	520	(2005)
Lifetime risk of maternal death (1 in N)	45	(2005)
Total maternal deaths	4,000	(2005)






INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %) 47 (2003) Wasting prevalence (moderate and severe, %)

5 (2003)


Complementary feeding rate (6-9 months, %) Low birthweight incidence (%)


84 (2008) 15 (2008)






























Adolescent birth rate (births per 1,000 women) 185 (2001) Unmet need for family planning (%) 18 (2003-04) Antenatal visits for woman (4 or more visits, %) (2003) 53 Intermittent preventive treatment for malaria (%) 43 (2008) C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) 2,5,1 (2003)Early initiation of breastfeeding (within 1 hr of birth, %) (2008)Postnatal visit for baby (within 2 days for home births, %)














WATER AND SANITATION







EQUITY

POLICIES

International Code of Marketing of Breastmilk Substitutes	Yes
New ORS formula and zinc for management of diarrhoea	Partial
Community treatment of pneumonia with antibiotics	Partial
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Partial
Midwives be authorised to administer a core set of life saving interventions	Yes
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	Partial

SYSTEMS

Financial Flows and Human Resources			
Per capita total expenditure on health (US\$)	39	(2007)	
General government expenditure on health as % of total government expenditure (%)	13	(2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	12	(2007)	
Density of health workers (per 10,000 population)	3.4	(2006)	
Official Development Assistance to child health per child (US\$)	14	(2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	d 33	(2007)	
National availability of Emergency Obstetric Car services (% of recommended minimum)	e 38	(2007)	


Mozambique




Myanmar

DEMOGRAPHICS

Total population (000)	49,563	(2008)
Total under-five population (000)	4,629	(2008)
Births (000)	1020	(2008)
Birth registration (%)	65	(2003)
Under-five mortality rate (per 1000 live births)	98	(2008)
Infant mortality rate (per 1000 live births)	71	(2008)
Neonatal mortality rate (per 1000 live births)	48	(2008)
Total under-five deaths (000)	98	(2008)
Maternal mortality ratio (per 100,000 live births)	380	(2005)
Lifetime risk of maternal death (1 in N)	110	(2005)
Total maternal deaths	3,700	(2005)






INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %) 41 (2003) Wasting prevalence (moderate and severe, %)


11 (2003)


Complementary feeding rate (6-9 months, %) Low birthweight incidence (%)


66 (2003)























Adolescent birth rate (births per 1,000 women)

Unmet need for family planning (%)

19 (2001)

Antenatal visits for woman (4 or more visits, %)

22 (2001)

Intermittent preventive treatment for malaria (%)


NA*


C-section rate (total, urban, rural; %)
(Minimum target is 5% and maximum target is 15%)


Early initiation of breastfeeding (within 1 hr of birth, %)


Postnatal visit for baby (within 2 days for home births, %)


^{*} Not applicable














WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	No
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	No
IMCI adapted to cover newborns 0-1 week of age	No
Costed implementation plan(s) for maternal, newborn and child health available	Partial
Midwives be authorised to administer a core set of life saving interventions	Partial
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	Yes

SYSTEMS

Financial Flows and Human Resources			
Per capita total expenditure on health (US\$)	21	(2007)	
General government expenditure on health as % of total government expenditure (%)	1	(2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	84	(2007)	
Density of health workers (per 10,000 population) 13	3.4	(2004)	
Official Development Assistance to child health per child (US\$)	2	(2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	4	(2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)) 		


Myanmar




Nepal

DEMOGRAPHICS

Total population (000)	28,810	(2008)
Total under-five population (000)	3,535	(2008)
Births (000)	732	(2008)
Birth registration (%)	35	(2006)
Under-five mortality rate (per 1000 live births)	51	(2008)
Infant mortality rate (per 1000 live births)	41	(2008)
Neonatal mortality rate (per 1000 live births)	31	(2008)
Total under-five deaths (000)	37	(2008)
Maternal mortality ratio (per 100,000 live births)	830	(2005)
Lifetime risk of maternal death (1 in N)	31	(2005)
Total maternal deaths	6,500	(2005)







INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %) 49 (2006) Wasting prevalence (moderate and severe, %) 13 (2006)


Complementary feeding rate (6-9 months, %) Low birthweight incidence (%)

75 (2006) 21 (2006)









Prevention of mother to child

2006

Percent HIV+ pregnant women receiving ARVs for PMTCT

2007

2008


25


transmission of HIV


10


8 6


4












Sub-national risk of malaria transmission

Adolescent birth rate (births per 1,000 women) 106 (2004) Unmet need for family planning (%) 25 (2006) Antenatal visits for woman (4 or more visits, %) 29 (2006) Intermittent preventive treatment for malaria (%) NA* C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) 3, 8, 2 (2006)


Early initiation of breastfeeding (within 1 hr of birth, %) 35 (2006) 2 (2006)


Postnatal visit for baby (within 2 days for home births, %)

















WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Yes
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	Yes
IMCI adapted to cover newborns 0-1 week of age	Partial
Costed implementation plan(s) for maternal, newborn and child health available	Partial
Midwives be authorised to administer a core set of life saving interventions	Partial
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	Partial

SYSTEMS

Financial Flows and Human Resources			
Per capita total expenditure on health (US\$)	53	(2007)	
General government expenditure on health as % of total government expenditure (%)	11	(2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	55	(2007)	
Density of health workers (per 10,000 population)	6.7	(2004)	
Official Development Assistance to child health per child (US\$)	5	(2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	16	(2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)			


Nepal




Niger

DEMOGRAPHICS

Total population (000)	14,704	(2008)
Total under-five population (000)	3,121	(2008)
Births (000)	791	(2008)
Birth registration (%)	32	(2006)
Under-five mortality rate (per 1000 live births)	167	(2008)
Infant mortality rate (per 1000 live births)	79	(2008)
Neonatal mortality rate (per 1000 live births)	34	(2008)
Total under-five deaths (000)	121	(2008)
Maternal mortality ratio (per 100,000 live births)	1,800	(2005)
Lifetime risk of maternal death (1 in N)	7	(2005)
Total maternal deaths	14,000	(2005)





INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %) 47 (2008) Wasting prevalence (moderate and severe, %)

12 (2008)


Complementary feeding rate (6-9 months, %) Low birthweight incidence (%)

66 (2008)


27 (2006)











Based on 2006 WHO reference population














Adolescent birth rate (births per 1,000 women) 199 (2004)

Unmet need for family planning (%) 16 (2006)

Antenatal visits for woman (4 or more visits, %) 15 (2006)


Intermittent preventive treatment for malaria (%) 0 (2006)


C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) 1, 5, 0 (2006)


Early initiation of breastfeeding (within 1 hr of birth, %) 38 (2008)


Postnatal visit for baby (within 2 days for home births, %) ---














WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Partial
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	Yes
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	Yes
Maternity protection in accordance with ILO Convention 183	Partial
Specific notification of maternal deaths	No

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	35 (2007)	
General government expenditure on health as % of total government expenditure (%)	12 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	45 (2007)	
Density of health workers (per 10,000 population)	1.6 (2008)	
Official Development Assistance to child health per child (US\$)	10 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	11 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)	68 (2000)	


Niger




Nigeria

DEMOGRAPHICS

Total population (000)	151,212	(2008)
Total under-five population (000)	25,020	(2008)
Births (000)	6,028	(2008)
Birth registration (%)	30	(2008)
Under-five mortality rate (per 1000 live births)	186	(2008)
Infant mortality rate (per 1000 live births)	96	(2008)
Neonatal mortality rate (per 1000 live births)	49	(2008)
Total under-five deaths (000)	1,077	(2008)
Maternal mortality ratio (per 100,000 live births) 1,100	(2005)
Lifetime risk of maternal death (1 in N)	18	(2005)
Total maternal deaths	59,000	(2005)







INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN NUTRITION


Stunting prevalence (moderate and severe, %) 41 (2008)

Wasting prevalence (moderate and severe, %) 14 (2008) Complementary feeding rate (6-9 months, %) Low birthweight incidence (%)


75 (2008) 14 (2003)











Based on 2006 WHO reference population


















Adolescent birth rate (births per 1,000 women) 123 (2006)
Unmet need for family planning (%) 20 (2008)
Antenatal visits for woman (4 or more visits, %) 45 (2008)
Intermittent preventive treatment for malaria (%) 5 (2008)
C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) 2, 4, 1 (2008)
Early initiation of breastfeeding (within 1 hr of birth, %) 38 (2008)
Postnatal visit for baby (within 2 days for home births, %) ---














WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Yes
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	Yes
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	Partial
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	No
Specific notification of maternal deaths	No

SYSTEMS

Financial Flows and Human Resources			
Per capita total expenditure on health (US\$)	131 (2007	')	
General government expenditure on health as % of total government expenditure (%)	7 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	72 (2007)	
Density of health workers (per 10,000 population)	20.1 (2008	3)	
Official Development Assistance to child health per child (US\$)	4 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	4 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)			


Nigeria




Pakistan

DEMOGRAPHICS

Total population (000)	176,952	(2008)
Total under-five population (000)	23,778	(2008)
Births (000)	5,337	(2008)
Birth registration (%)		
Under-five mortality rate (per 1000 live births)	89	(2008)
Infant mortality rate (per 1000 live births)	72	(2008)
Neonatal mortality rate (per 1000 live births)	53	(2008)
Total under-five deaths (000)	465	(2008)
Maternal mortality ratio (per 100,000 live births) 320	(2005)
Lifetime risk of maternal death (1 in N)	74	(2005)
Total maternal deaths	15,000	(2005)







INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %) 42 (2001-2002) Wasting prevalence (moderate and severe, %) 14 (2001-2002)

Low birthweight incidence (%)


Complementary feeding rate (6-9 months, %) 36 (2006-2007) 32 (2006-2007)











Based on 2006 WHO reference population














 $\begin{tabular}{lll} Adolescent birth rate (births per 1,000 women) & 20 & (2005) \\ Unmet need for family planning (%) & 25 & (2006-07) \\ Antenatal visits for woman (4 or more visits, %) & 28 & (2006-07) \\ Intermittent preventive treatment for malaria (%) & NA* \\ \end{tabular}$


C-section rate (total, urban, rural; %)

(Minimum target is 5% and maximum target is 15%) 7,13,**5** (2006-07)


Early initiation of breastfeeding (within 1 hr of birth, %) 29 (2006-07)


Postnatal visit for baby (within 2 days for home births, %) ---


^{*} Not applicable














WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Partial
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	Yes
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	Partial
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	No

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	64 (2007)	
General government expenditure on health as % of total government expenditure (%)	4 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	57 (2007)	
Density of health workers (per 10,000 population)	11.6 (2007)	
Official Development Assistance to child health per child (US\$)	5 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	9 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)		


Pakistan

Papua New Guinea

DEMOGRAPHICS

Total population (000)	6,577	(2008)
Total under-five population (000)	950	(2008)
Births (000)	207	(2008)
Birth registration (%)		
Under-five mortality rate (per 1000 live births)	69	(2008)
Infant mortality rate (per 1000 live births)	53	(2008)
Neonatal mortality rate (per 1000 live births)	26	(2008)
Total under-five deaths (000)	14	(2008)
Maternal mortality ratio (per 100,000 live births)	470	(2005)
Lifetime risk of maternal death (1 in N)	55	(2005)
Total maternal deaths	820	(2005)







INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %) 43 (2005) Wasting prevalence (moderate and severe, %) 5 (2005)

Complementary feeding rate (6-9 months, %) Low birthweight incidence (%)


76 (2006)









CHILD HEALTH















*Only includes children under 3 years of age


Adolescent birth rate (births per 1,000 women) 70 (2000)


Unmet need for family planning (%) --
Antenatal visits for woman (4 or more visits, %) 55 (2006)


Intermittent preventive treatment for malaria (%) --
C-section rate (total, urban, rural; %)
(Minimum target is 5% and maximum target is 15%) ---, ---, --
Early initiation of breastfeeding (within 1 hr of birth, %) --
Postnatal visit for baby (within 2 days for home births, %) ---













WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Partial
New ORS formula and zinc for management of diarrhoea	No
Community treatment of pneumonia with antibiotics	No
IMCI adapted to cover newborns 0-1 week of age	Partial
Costed implementation plan(s) for maternal, newborn and child health available	Partial
Midwives be authorised to administer a core set of life saving interventions	Partial
Maternity protection in accordance with ILO Convention 183	Partial
Specific notification of maternal deaths	Yes

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	65 (2007)	
General government expenditure on health as % of total government expenditure (%)	7 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	8 (2007)	
Density of health workers (per 10,000 population)	5.8 (2000)	
Official Development Assistance to child health per child (US\$)	11 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	24 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)		


Papua New Guinea




Peru

DEMOGRAPHICS

Total population (000)	28,837	(2008)
Total under-five population (000)	2,975	(2008)
Births (000)	609	(2008)
Birth registration (%)	93	(2006)
Under-five mortality rate (per 1000 live births)	24	(2008)
Infant mortality rate (per 1000 live births)	22	(2008)
Neonatal mortality rate (per 1000 live births)	13	(2008)
Total under-five deaths (000)	15	(2008)
Maternal mortality ratio (per 100,000 live births)	240	(2005)
Lifetime risk of maternal death (1 in N)	140	(2005)
Total maternal deaths	1,500	(2005)






INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %) 30 (2004-2006)


Wasting prevalence (moderate and severe, %) 1 (2004-2006)


Underweight prevalence Percent children < 5 years underweight for age* 100 80 60 40 20 6 0 2004-2006 ased on 2006 WHO reference population


Complementary feeding rate (6-9 months, %) Low birthweight incidence (%) 8 (2007)





















Adolescent birth rate (births per 1,000 women) 59 (2004)
Unmet need for family planning (%) 8 (2004-06)
Antenatal visits for woman (4 or more visits, %) 87 (2004-06)
Intermittent preventive treatment for malaria (%) NA*


C-section rate (total, urban, **rural**; %) (Minimum target is 5% and maximum target is 15%) 16, 24, **6** (2004-06)


Early initiation of breastfeeding (within 1 hr of birth, %) 48 (2004-06)


Postnatal visit for baby (within 2 d1ays for home births, %) ---


^{*} Not applicable














WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Yes
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	Partial
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	Partial
Maternity protection in accordance with ILO Convention 183	Partial
Specific notification of maternal deaths	Yes

SYSTEMS

٠.			
	Financial Flows and Human Resources		
	Per capita total expenditure on health (US\$)	327 (2007)	
	General government expenditure on health as % of total government expenditure (%)	16 (2007)	
	Out-of-pocket expenditure as % of total expenditure on health (%)	31 (2007)	
	Density of health workers (per 10,000 population)	18.4 (1999)	
	Official Development Assistance to child health per child (US\$)	3 (2007)	
	Official Development Assistance to maternal and neonatal health per live birth (US\$)	11 (2007)	
	National availability of Emergency Obstetric Care services (% of recommended minimum)		


Peru




Philippines

DEMOGRAPHICS

Total population (000)	90,348	(2008)
Total under-five population (000)	10,701	(2008)
Births (000)	2,236	(2008)
Birth registration (%)	83	(2000)
Under-five mortality rate (per 1000 live births)	32	(2008)
Infant mortality rate (per 1000 live births)	26	(2008)
Neonatal mortality rate (per 1000 live births)	15	(2008)
Total under-five deaths (000)	73	(2008)
Maternal mortality ratio (per 100,000 live births)	230	(2005)
Lifetime risk of maternal death (1 in N)	140	(2005)
Total maternal deaths	4,600	(2005)





INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %) 34 (2003) Wasting prevalence (moderate and severe, %)


6 (2003)


Complementary feeding rate (6-9 months, %) Low birthweight incidence (%)

58 (2008)


20 (2003)











CHILD HEALTH















Sub-national risk of malaria transmission

Adolescent birth rate (births per 1,000 women) 54 (2007)
Unmet need for family planning (%) 17 (2003)
Antenatal visits for woman (4 or more visits, %) 70 (2003)
Intermittent preventive treatment for malaria (%) NA*


C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) 7, 10, **5** (2003) Early initiation of breastfeeding (within 1 hr of birth, %) 54 (2003)


Postnatal visit for baby (within 2 days for home births, %) ---


* Not applicable














WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Yes
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	No
IMCI adapted to cover newborns 0-1 week of age	No
Costed implementation plan(s) for maternal, newborn and child health available	No
Midwives be authorised to administer a core set of life saving interventions	No
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	Partial

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	130 (2007)	
General government expenditure on health as % of total government expenditure (%)	7 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	55 (2007)	
Density of health workers (per 10,000 population)	72.7 (2002)	
Official Development Assistance to child health per child (US\$)	2 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	5 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)		


Philippines




Rwanda

DEMOGRAPHICS

Total population (000)	9,721	(2008)
Total under-five population (000)	1,646	(2008)
Births (000)	403	(2008)
Birth registration (%)	82	(2005)
Under-five mortality rate (per 1000 live births)	112	(2008)
Infant mortality rate (per 1000 live births)	72	(2008)
Neonatal mortality rate (per 1000 live births)	35	(2008)
Total under-five deaths (000)	41	(2008)
Maternal mortality ratio (per 100,000 live births)	1,300	(2005)
Lifetime risk of maternal death (1 in N)	16	(2005)
Total maternal deaths	4,700	(2005)








INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


NUTRITION


Stunting prevalence (moderate and severe, %) 51 (2005) Wasting prevalence (moderate and severe, %) 5 (2005) Complementary feeding rate (6-9 months, %) Low birthweight incidence (%) 69 (2005) 6 (2005)


























(2006) Adolescent birth rate (births per 1,000 women) 40 Unmet need for family planning (%) 38 (2005)Antenatal visits for woman (4 or more visits, %) 24 (2007-08) Intermittent preventive treatment for malaria (%) 17 (2007-08) C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) 3, 8, 2 (2005)Early initiation of breastfeeding (within 1 hr of birth, %) (2005)Postnatal visit for baby (within 2 days for home births, %)














WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	No
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	Yes
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	Yes
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	No
L	

SYSTEMS

Financial Flows and Human Resources			
Per capita total expenditure on health (US\$)	90 (2007)		
General government expenditure on health as % of total government expenditure (%)	20 (2007)		
Out-of-pocket expenditure as % of total expenditure on health (%)	25 (2007)		
Density of health workers (per 10,000 population)	4.7 (2005)		
Official Development Assistance to child health per child (US\$)	15 (2007)		
Official Development Assistance to maternal and neonatal health per live birth (US\$)	15 (2007)		
National availability of Emergency Obstetric Care services (% of recommended minimum)			


Rwanda




Senegal

DEMOGRAPHICS

Total population (000)	12,211	(2008)
Total under-five population (000)	2,046	(2008)
Births (000)	470	(2008)
Birth registration (%)	55	(2005)
Under-five mortality rate (per 1000 live births)	108	(2008)
Infant mortality rate (per 1000 live births)	57	(2008)
Neonatal mortality rate (per 1000 live births)	34	(2008)
Total under-five deaths (000)	49	(2008)
Maternal mortality ratio (per 100,000 live births)	980	(2005)
Lifetime risk of maternal death (1 in N)	21	(2005)
Total maternal deaths	4,100	(2005)





INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %) 19 (2005) Wasting prevalence (moderate and severe, %)


9 (2005)


Complementary feeding rate (6-9 months, %) Low birthweight incidence (%)


61 (2005)


19 (2005)



























Adolescent birth rate (births per 1,000 women) 96 (2007)Unmet need for family planning (%) 32 (2005)Antenatal visits for woman (4 or more visits, %) 40 (2005)Intermittent preventive treatment for malaria (%) 52 (2008-09) C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) 3, 7, 1 (2005)Early initiation of breastfeeding (within 1 hr of birth, %) (2005)Postnatal visit for baby (within 2 days for home births, %) ---














WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Partial
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	Yes
IMCI adapted to cover newborns 0-1 week of age	No
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	Yes
Maternity protection in accordance with ILO Convention 183	Partial
Specific notification of maternal deaths	No

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	99 (2007)	
General government expenditure on health as % of total government expenditure (%)	12 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	35 (2007)	
Density of health workers (per 10,000 population)	4.8 (2008)	
Official Development Assistance to child health per child (US\$)	8 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	10 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)	38 (2000)	


Senegal




Sierra Leone

DEMOGRAPHICS

Total population (000)	5,560	(2008)
Total under-five population (000)	947	(2008)
Births (000)	223	(2008)
Birth registration (%)	51	(2008)
Under-five mortality rate (per 1000 live births)	194	(2008)
Infant mortality rate (per 1000 live births)	123	(2008)
Neonatal mortality rate (per 1000 live births)	45	(2008)
Total under-five deaths (000)	43	(2008)
Maternal mortality ratio (per 100,000 live births)	2,100	(2005)
Lifetime risk of maternal death (1 in N)	8	(2005)
Total maternal deaths	5,400	(2005)







INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %) 36 (2008) Wasting prevalence (moderate and severe, %) 10 (2008)

Complementary feeding rate (6-9 months, %) Low birthweight incidence (%)


73 (2008) 24 (2005)











Based on 2006 WHO reference population

















Adolescent birth rate (births per 1,000 women) 143 (2006)
Unmet need for family planning (%) 28 (2008)
Antenatal visits for woman (4 or more visits, %) 56 (2008)
Intermittent preventive treatment for malaria (%) 10 (2008)
C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) 2, 3, 1 (2008)
Early initiation of breastfeeding (within 1 hr of birth, %) 51 (2008)
Postnatal visit for baby (within 2 days for home births, %) ---














WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	No
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	Yes
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Partial
Midwives be authorised to administer a core set of life saving interventions	Partial
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	Partial

SYSTEMS

Financial Flows and Human Resources			
Per capita total expenditure on health (US\$)	32 (2007)		
General government expenditure on health as % of total government expenditure (%)	8 (2007)		
Out-of-pocket expenditure as % of total expenditure on health (%)	40 (2007)		
Density of health workers (per 10,000 population)	1.9 (2008)		
Official Development Assistance to child health per child (US\$)	14 (2007)		
Official Development Assistance to maternal and neonatal health per live birth (US\$)	d 35 (2007)		
National availability of Emergency Obstetric Care services (% of recommended minimum)	e 24 (2008)		


Sierra Leone




Somalia

DEMOGRAPHICS

Total population (000)	8,926	(2008)
Total under-five population (000)	1,611	(2008)
Births (000)	395	(2008)
Birth registration (%)	3	(2006)
Under-five mortality rate (per 1000 live births)	200	(2008)
Infant mortality rate (per 1000 live births)	119	(2008)
Neonatal mortality rate (per 1000 live births)	61	(2008)
Total under-five deaths (000)	76	(2008)
Maternal mortality ratio (per 100,000 live births)	1,400	(2005)
Lifetime risk of maternal death (1 in N)	12	(2005)
Total maternal deaths	5,200	(2005)





INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %) 42 (2006) Wasting prevalence (moderate and severe, %)


13 (2006)


Low birthweight incidence (%)

Complementary feeding rate (6-9 months, %)


15 (2006)











Based on 2006 WHO reference population






























WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	No
New ORS formula and zinc for management of diarrhoea	Partial
Community treatment of pneumonia with antibiotics	No
IMCI adapted to cover newborns 0-1 week of age	No
Costed implementation plan(s) for maternal, newborn and child health available	No
Midwives be authorised to administer a core set of life saving interventions	No
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	No

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)		
General government expenditure on health as % of total government expenditure (%)		
Out-of-pocket expenditure as % of total expenditure on health (%)		
Density of health workers (per 10,000 population)	1.5 (2006)	
Official Development Assistance to child health per child (US\$)	10 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	16 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)	56 (2005)	


Somalia




South Africa

DEMOGRAPHICS

Total population (000)	49,668	(2008)
Total under-five population (000)	5,200	(2008)
Births (000)	1,091	(2008)
Birth registration (%)	78	(2006)
Under-five mortality rate (per 1000 live births)	67	(2008)
Infant mortality rate (per 1000 live births)	48	(2008)
Neonatal mortality rate (per 1000 live births)	20	(2008)
Total under-five deaths (000)	73	(2008)
Maternal mortality ratio (per 100,000 live births)	400	(2005)
Lifetime risk of maternal death (1 in N)	110	(2005)
Total maternal deaths	4,300	(2005)






INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %) 33 (2003) Wasting prevalence (moderate and severe, %)


6 (2003)

Complementary feeding rate (6-9 months, %) Low birthweight incidence (%)


49 (2003)









Based on 2006 WHO reference population


















Adolescent birth rate (births per 1,000 women) 54 (2007)
Unmet need for family planning (%) 15 (1998)
Antenatal visits for woman (4 or more visits, %) 56 (2003)
Intermittent preventive treatment for malaria (%) NA*


C-section rate (total, urban, **rural**; %)
(Minimum target is 5% and maximum target is 15%) 21, 24, **15** (2003)
Early initiation of breastfeeding (within 1 hr of birth, %) 61 (2003)


Postnatal visit for baby (within 2 days for home births, %)

















WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Partial
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	No
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	Yes
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	Yes

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	819 (2007)	
General government expenditure on health as % of total government expenditure (%)	11 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	17 (2007)	
Density of health workers (per 10,000 population)	48.5 (2004)	
Official Development Assistance to child health per child (US\$)	4 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	3 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)		


South Africa




Sudan

DEMOGRAPHICS

Total population (000)	41,348	(2008)
Total under-five population (000)	5,836	(2008)
Births (000)	1,296	(2008)
Birth registration (%)	33	(2006)
Under-five mortality rate (per 1000 live births)	109	(2008)
Infant mortality rate (per 1000 live births)	70	(2008)
Neonatal mortality rate (per 1000 live births)	41	(2008)
Total under-five deaths (000)	138	(2008)
Maternal mortality ratio (per 100,000 live births)	450	(2005)
Lifetime risk of maternal death (1 in N)	53	(2005)
Total maternal deaths	5,300	(2005)






INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %) 40 (2006) Wasting prevalence (moderate and severe, %)


16 (2006)

Complementary feeding rate (6-9 months, %) Low birthweight incidence (%)


56 (2006) 31 (1999)











Based on 2006 WHO reference population














Adolescent birth rate (births per 1,000 women) 72 (1997)


Unmet need for family planning (%) 26 (1992-93)


Antenatal visits for woman (4 or more visits, %) --
Intermittent preventive treatment for malaria (%) --
C-section rate (total, urban, rural; %)
(Minimum target is 5% and maximum target is 15%) 20,19,24 (1992-93)


Early initiation of breastfeeding (within 1 hr of birth, %) ---


Postnatal visit for baby (within 2 days for home births, %)













WATER AND SANITATION







EQUITY

POLICIES

International Code of Marketing of Breastmilk Substitutes	Partial
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	Yes
IMCI adapted to cover newborns 0-1 week of age	No
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	No
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	Partial

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	71 (2007)	
General government expenditure on health as % of total government expenditure (%)	6 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	63 (2007)	
Density of health workers (per 10,000 population)	12.0 (2006)	
Official Development Assistance to child health per child (US\$)	11 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	14 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)	35 (2005)	


Sudan




Swaziland

DEMOGRAPHICS

Total population (000)	1,168	(2008)
Total under-five population (000)	159	(2008)
Births (000)	35	(2008)
Birth registration (%)	30	(2006-07)
Under-five mortality rate (per 1000 live births)	83	(2008)
Infant mortality rate (per 1000 live births)	59	(2008)
Neonatal mortality rate (per 1000 live births)	18	(2008)
Total under-five deaths (000)	3	(2008)
Maternal mortality ratio (per 100,000 live births	390	(2005)
Lifetime risk of maternal death (1 in N)	120	(2005)
Total maternal deaths	120	(2005)







INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %) 29 (2006-2007)

Wasting prevalence (moderate and severe, %) 3 (2006-2007)



Complementary feeding rate (6-9 months, %) 77 (2006-2007) Low birthweight incidence (%) 9 (2006-2007)








Prevention of mother to child


Percent HIV+ pregnant women receiving ARVs for PMTCT


transmission of HIV















Adolescent birth rate (births per 1,000 women) 111 (2004)
Unmet need for family planning (%) 24 (2006-07)
Antenatal visits for woman (4 or more visits, %) 79 (2006-07)
Intermittent preventive treatment for malaria (%) 1 (2006-07)
C-section rate (total, urban, rural; %)

C-section rate (total, urban, **rural**; %) (Minimum target is 5% and maximum target is 15%) 8, 8, **8** (2006-07)


Early initiation of breastfeeding (within 1 hr of birth, %) 67 (2006-07)


Postnatal visit for baby (within 2 days for home births, %) ---







Coverage along the continuum of care









WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Partial
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	No
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	No
Midwives be authorised to administer a core set of life saving interventions	Yes
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	Yes

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	287 (2007)	
General government expenditure on health as % of total government expenditure (%)	9 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	16 (2007)	
Density of health workers (per 10,000 population)	64.6 (2004)	
Official Development Assistance to child health per child (US\$)	16 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	17 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)		


Swaziland




Tajikistan

DEMOGRAPHICS

Total population (000)	6,836	(2008)
Total under-five population (000)	871	(2008)
Births (000)	193	(2008)
Birth registration (%)	88	(2005)
Under-five mortality rate (per 1000 live births)	64	(2008)
Infant mortality rate (per 1000 live births)	54	(2008)
Neonatal mortality rate (per 1000 live births)	22	(2008)
Total under-five deaths (000)	12	(2008)
Maternal mortality ratio (per 100,000 live births)	170	(2005)
Lifetime risk of maternal death (1 in N)	160	(2005)
Total maternal deaths	320	(2005)





INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %) Wasting prevalence (moderate and severe, %)


39 (2007) 7 (2007)


Complementary feeding rate (6-9 months, %) Low birthweight incidence (%)

15 (2005)

10 (2005)











Based on 2006 WHO reference population


CHILD HEALTH













Prevention of mother to child

transmission of HIV




Postnatal visit for baby (within 2 days for home births, %) --

















WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	No
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	Yes
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Partial
Midwives be authorised to administer a core set of life saving interventions	Partial
Maternity protection in accordance with ILO Convention 183	Partial
Specific notification of maternal deaths	Yes

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	93 (2007)	
General government expenditure on health as % of total government expenditure (%)	4 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	74 (2007)	
Density of health workers (per 10,000 population)	70.4 (2006)	
Official Development Assistance to child health per child (US\$)	6 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	10 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)	86 (2005)	


Tajikistan




Tanzania, United Republic of

DEMOGRAPHICS

Total population (000)	42,484	(2008)
	,	` ,
Total under-five population (000)	7,566	(2008)
Births (000)	1,771	(2008)
Birth registration (%)	8	(2004-05)
Under-five mortality rate (per 1000 live bir	ths) 104	(2008)
Infant mortality rate (per 1000 live births)	67	(2008)
Neonatal mortality rate (per 1000 live birth	ns) 33	(2008)
Total under-five deaths (000)	175	(2008)
Maternal mortality ratio (per 100,000 live bi	irths) 950	(2005)
Lifetime risk of maternal death (1 in N)	24	(2005)
Total maternal deaths	13,000	(2005)






INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %) 44 (2004-2005)


Wasting prevalence (moderate and severe, %) 4 (2004-2005)


Underweight prevalence Percent children < 5 years underweight for age*





Complementary feeding rate (6-9 months, %) 91 (2004-2005) Low birthweight incidence (%) 10 (2004-2005)




















Adolescent birth rate (births per 1,000 women) 139 (2003)

Unmet need for family planning (%) 22 (2004-05)

Antenatal visits for woman (4 or more visits, %) 62 (2004-05)


Intermittent preventive treatment for malaria (%) 30 (2007-08)


C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) 3,8,2 (2004-05)

Early initiation of breastfeeding (within 1 hr of birth, %) 67 (2004-05)


Postnatal visit for baby (within 2 days for home births, %)














WATER AND SANITATION









POLICIES

International Code of Marketing of Breastmilk Substitutes	Yes
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	No
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	Yes
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	Yes

SYSTEMS


Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	63 (2007)	
General government expenditure on health as % of total government expenditure (%)	18 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	26 (2007)	
Density of health workers (per 10,000 population)	2.5 (2006)	
Official Development Assistance to child health per child (US\$)	11 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	19 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)	21 (2005)	


Tanzania, United Republic of



DEMOGRAPHICS

Total population (000)	6,459	(2008)
Total under-five population (000)	947	(2008)
Births (000)	213	(2008)
Birth registration (%)	78	(2006)
Under-five mortality rate (per 1000 live births)	98	(2008)
Infant mortality rate (per 1000 live births)	64	(2008)
Neonatal mortality rate (per 1000 live births)	33	(2008)
Total under-five deaths (000)	20	(2008)
Maternal mortality ratio (per 100,000 live births)	510	(2005)
Lifetime risk of maternal death (1 in N)	38	(2005)
Total maternal deaths	1,200	(2005)






INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %) 27 (2008) Wasting prevalence (moderate and severe, %)


6 (2008)

Complementary feeding rate (6-9 months, %) Low birthweight incidence (%)


70 (2008) 12 (2006)











Based on 2006 WHO reference population

















Adolescent birth rate (births per 1,000 women)	89	(1996)
Unmet need for family planning (%)	32	(1998)
Antenatal visits for woman (4 or more visits, %)	46	(1998)
Intermittent preventive treatment for malaria (%)	18	(2006)
C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%)	2, 5, 1	(1998)
Early initiation of breastfeeding (within 1 hr of birth, %)	53	(2008)


Postnatal visit for baby (within 2 days for home births, %)














WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Partial
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	Yes
IMCI adapted to cover newborns 0-1 week of age	No
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	Yes
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	Yes

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	68 (2007)	
General government expenditure on health as % of total government expenditure (%)	8 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	63 (2007)	
Density of health workers (per 10,000 population)	3.2 (2008)	
Official Development Assistance to child health per child (US\$)	8 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	11 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)		


Togo




Turkmenistan

DEMOGRAPHICS

Total population (000)	5,044	(2008)
Total under-five population (000)	518	(2008)
Births (000)	111	(2008)
Birth registration (%)	96	(2006)
Under-five mortality rate (per 1000 live births)	48	(2008)
Infant mortality rate (per 1000 live births)	43	(2008)
Neonatal mortality rate (per 1000 live births)	21	(2008)
Total under-five deaths (000)	5	(2008)
Maternal mortality ratio (per 100,000 live births)	130	(2005)
Lifetime risk of maternal death (1 in N)	290	(2005)
Total maternal deaths	140	(2005)






INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN

NUTRITION


Stunting prevalence (moderate and severe, %) 19 (2006) Wasting prevalence (moderate and severe, %) 7 (2006) Complementary feeding rate (6-9 months, %) 54 (2006) Low birthweight incidence (%) 4 (2006)










*Based on 2006 WHO reference population














Adolescent birth rate (births per 1,000 women)

21 (2006)

Unmet need for family planning (%)

Antenatal visits for woman (4 or more visits, %)

Intermittent preventive treatment for malaria (%)

C-section rate (total, urban, rural; %)

(Missingura tyract in 58% and proving to pract in 158%)

21 (2006)


NA*


(Minimum target is 5% and maximum target is 15%) 3, 4, 2 (2000)


Early initiation of breastfeeding (within 1 hr of birth, %) 60 (2006)


Postnatal visit for baby (within 2 days for home births, %) -

















WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Yes
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	Partial
IMCI adapted to cover newborns 0-1 week of age	No
Costed implementation plan(s) for maternal, newborn and child health available	Partial
Midwives be authorised to administer a core set of life saving interventions	Partial
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	No

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	153 (2007)	
General government expenditure on health as % of total government expenditure (%)	10 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	48 (2007)	
Density of health workers (per 10,000 population)	69.6 (2007)	
Official Development Assistance to child health per child (US\$)	3 (2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	6 (2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)		


Turkmenistan




Uganda

DEMOGRAPHICS

Total population (000)	31,657	(2008)
Total under-five population (000)	6,182	(2008)
Births (000)	1,466	(2008)
Birth registration (%)	21	(2006)
Under-five mortality rate (per 1000 live births)	135	(2008)
Infant mortality rate (per 1000 live births)	85	(2008)
Neonatal mortality rate (per 1000 live births)	31	(2008)
Total under-five deaths (000)	190	(2008)
Maternal mortality ratio (per 100,000 live births)	550	(2005)
Lifetime risk of maternal death (1 in N)	25	(2005)
Total maternal deaths	8,100	(2005)





INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %)


38 (2006) 6 (2006)


Complementary feeding rate (6-9 months, %) 80 (2006)


Low birthweight incidence (%)


14 (2006)


























Adolescent birth rate (births per 1,000 women) 159 (2004)
Unmet need for family planning (%) 41 (2006)
Antenatal visits for woman (4 or more visits, %) 47 (2006)
Intermittent preventive treatment for malaria (%) 16 (2006)
C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) 3, 9, 2 (2006)
Early initiation of breastfeeding (within 1 hr of birth, %) 42 (2006)
Postnatal visit for baby (within 2 days for home births, %) ---














WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Yes
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	Yes
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	Partial
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	Partial

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	74 (2007)	
General government expenditure on health as % of total government expenditure (%)	10 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	38 (2007)	
Density of health workers (per 10,000 population)	14.3 (2005)	
Official Development Assistance to child health per child (US\$)	13 (2007)	
Official Development Assistance to maternal ar neonatal health per live birth (US\$)	nd 12 (2007)	
National availability of Emergency Obstetric Ca services (% of recommended minimum)	re 34 (2002-2003)	


Uganda




Yemen

DEMOGRAPHICS

Total population (000)	22,917	(2008)
Total under-five population (000)	3,733	(2008)
Births (000)	846	(2008)
Birth registration (%)	22	(2006)
Under-five mortality rate (per 1000 live births)	69	(2008)
Infant mortality rate (per 1000 live births)	53	(2008)
Neonatal mortality rate (per 1000 live births)	32	(2008)
Total under-five deaths (000)	57	(2008)
Maternal mortality ratio (per 100,000 live births)	430	(2005)
Lifetime risk of maternal death (1 in N)	39	(2005)
Total maternal deaths	3,600	(2005)







INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN


Stunting prevalence (moderate and severe, %) 58 (2003) Wasting prevalence (moderate and severe, %) 14 (2003)


Complementary feeding rate (6-9 months, %) Low birthweight incidence (%)


76 (2003) 32 (1997)























Adolescent birth rate (births per 1,000 women) 80 (2005)
Unmet need for family planning (%) 39 (1997)
Antenatal visits for woman (4 or more visits, %) 11 (1997)
Intermittent preventive treatment for malaria (%) NA*

C-section rate (total, urban, rural; %)
(Minimum target is 5% and maximum target is 15%) 1, 2, 1 (1997)

Early initiation of breastfeeding (within 1 hr of birth, %) 30 (2006)


Postnatal visit for baby (within 2 days for home births, %)

















WATER AND SANITATION





EQUITY



POLICIES

_		
	International Code of Marketing of Breastmilk Substitutes	Yes
	New ORS formula and zinc for management of diarrhoea	Yes
	Community treatment of pneumonia with antibiotics	Yes
	IMCI adapted to cover newborns 0-1 week of age	No
	Costed implementation plan(s) for maternal, newborn and child health available	Yes
	Midwives be authorised to administer a core set of life saving interventions	Yes
	Maternity protection in accordance with ILO Convention 183	No
	Specific notification of maternal deaths	No
_		

SYSTEMS

Financial Flows and Human Resources		
Per capita total expenditure on health (US\$)	104 (2007)	
General government expenditure on health as % of total government expenditure (%)	5 (2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	59 (2007)	
Density of health workers (per 10,000 population)	9.9 (2004)	
Official Development Assistance to child health per child (US\$)	5 (2007)	
Official Development Assistance to maternal an neonatal health per live birth (US\$)	d 18 (2007)	
National availability of Emergency Obstetric Cal services (% of recommended minimum)	re 14 (2004-2005)	


Yemen




Zambia

DEMOGRAPHICS

Total population (000)	12,620	(2008)
Total under-five population (000)	2,282	(2008)
Births (000)	542	(2008)
Birth registration (%)	10	(2000)
Under-five mortality rate (per 1000 live births)	148	(2008)
Infant mortality rate (per 1000 live births)	92	(2008)
Neonatal mortality rate (per 1000 live births)	36	(2008)
Total under-five deaths (000)	77	(2008)
Maternal mortality ratio (per 100,000 live births)	830	(2005)
Lifetime risk of maternal death (1 in N)	27	(2005)
Total maternal deaths	3,900	(2005)





INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN

Stunting prevalence (moderate and severe, %) 45 (2007) Wasting prevalence (moderate and severe, %)


5 (2007)


Low birthweight incidence (%)


Complementary feeding rate (6-9 months, %)

93 (2007)


11 (2007)











Based on 2006 WHO reference population

















MATERNAL AND NEWBORN HEALTH


Adolescent birth rate (births per 1,000 women)	151	(2005)
Unmet need for family planning (%)	27	(2007)
Antenatal visits for woman (4 or more visits, %)	60	(2007)
Intermittent preventive treatment for malaria (%)	60	(2008)
C-section rate (total, urban, rural ; %) (Minimum target is 5% and maximum target is 15%)	3,6, 2	(2007)
Early initiation of breastfeeding (within 1 hr of birth, %)	57	(2007)
Postnatal visit for haby (within 2 days for home births %)		














WATER AND SANITATION





EQUITY



POLICIES

International Code of Marketing of Breastmilk Substitutes	Partial
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	Partial
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Partial
Midwives be authorised to administer a core set of life saving interventions	Yes
Maternity protection in accordance with ILO Convention 183	Partial
Specific notification of maternal deaths	Yes

SYSTEMS

Financial Flows and Human Resources			
Per capita total expenditure on health (US\$)	79	(2007)	
General government expenditure on health as % of total government expenditure (%)	15	(2007)	
Out-of-pocket expenditure as % of total expenditure on health (%)	29	(2007)	
Density of health workers (per 10,000 population)	7.7	(2006)	
Official Development Assistance to child health per child (US\$)	27	(2007)	
Official Development Assistance to maternal and neonatal health per live birth (US\$)	28	(2007)	
National availability of Emergency Obstetric Care services (% of recommended minimum)	41	(2004-2005)	


Zambia




Zimbabwe

DEMOGRAPHICS

Total population (000)	12,463	(2008)
Total under-five population (000)	1,707	(2008)
Births (000)	378	(2008)
Birth registration (%)	74	(2005-06)
Under-five mortality rate (per 1000 live births	s) 96	(2008)
Infant mortality rate (per 1000 live births)	62	(2008)
Neonatal mortality rate (per 1000 live births)	28	(2008)
Total under-five deaths (000)	36	(2008)
Maternal mortality ratio (per 100,000 live birth	s) 880	(2005)
Lifetime risk of maternal death (1 in N)	43	(2005)
Total maternal deaths	3,400	(2005)








INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN

Stunting prevalence (moderate and severe, %) 33 (2005-2006) Wasting prevalence (moderate and severe, %) 7 (2005-2006)


Complementary feeding rate (6-9 months, %) 79 (2005-2006) Low birthweight incidence (%) 11 (2005-2006)











CHILD HEALTH














MATERNAL AND NEWBORN HEALTH

Adolescent birth rate (births per 1,000 women) 101 (2003)

Unmet need for family planning (%) 13 (2005-06)

Antenatal visits for woman (4 or more visits, %) 71 (2005-06)


Intermittent preventive treatment for malaria (%) 6 (2005-06)


C-section rate (total, urban, rural; %) (Minimum target is 5% and maximum target is 15%) 5,9,3 (2005-06)


Early initiation of breastfeeding (within 1 hr of birth, %) 69 (2005-06)


Postnatal visit for baby (within 2 days for home births, %)














WATER AND SANITATION









POLICIES

International Code of Marketing of Breastmilk Substitutes	Yes
New ORS formula and zinc for management of diarrhoea	Yes
Community treatment of pneumonia with antibiotics	No
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Yes
Midwives be authorised to administer a core set of life saving interventions	Partial
Maternity protection in accordance with ILO Convention 183	No
Specific notification of maternal deaths	Yes

SYSTEMS

Financial Flows and Human Resources			
Per capita total expenditure on health (US\$)	20 (2007)		
General government expenditure on health as % of total government expenditure (%)	9 (2007)		
Out-of-pocket expenditure as % of total expenditure on health (%)	27 (2007)		
Density of health workers (per 10,000 population)	8.8 (2004)		
Official Development Assistance to child health per child (US\$)	18 (2007)		
Official Development Assistance to maternal and neonatal health per live birth (US\$)	21 (2007)		
National availability of Emergency Obstetric Care services (% of recommended minimum)			

Zimbabwe

Annex A Country profile indicators and data sources

Indicator		Data source	Global database
Demographics			
Demographics	Total population	United Nations Population Division	United Nations Population Division
	Total children under-five population	United Nations Population Division	United Nations Population Division
	Total births	United Nations Population Division	United Nations Population Division
	Birth registration	Multiple Indicator Cluster Surveys, Demographic and Health Surveys	United Nations Children's Fund
Child mortality	Under-five mortality rate	Inter-agency Group for Child Mortality Estimation (United Nations Children's Fund, World Health Organization, United Nations Population Division, World Bank)	Inter-agency Group for Child Mortality Estimation (United Nations Children's Fund, World Health Organization, United Nations Population Division, World Bank)
	Infant mortality rate	Inter-agency Group for Child Mortality Estimation (United Nations Children's Fund, World Health Organization, United Nations Population Division, World Bank)	Inter-agency Group for Child Mortality Estimation (United Nations Children's Fund, World Health Organization, United Nations Population Division, World Bank)
	Neonatal mortality rate	World Health Organization	World Health Organization
	Total deaths for children under age 5	Inter-agency Group for Child Mortality Estimation (United Nations Children's Fund, World Health Organization, United Nations Population Division, World Bank)	Inter-agency Group for Child Mortality Estimation (United Nations Children's Fund, World Health Organization, United Nations Population Division, World Bank)
	Cause of death of children under age 5	World Health Organization/Child Health Epidemiology Reference Group	World Health Organization/Child Health Epidemiology Reference Group
	Causes of neonatal deaths	World Health Organization/Child Health Epidemiology Reference Group	World Health Organization/Child Health Epidemiology Reference Group
Maternal mortality	Maternal mortality ratio (adjusted)	World Health Organization/United Nations Children's Fund/ United Nations Population Fund/World Bank	World Health Organization/United Nations Children's Fund/United Nations Population Fund/World Bank
	Lifetime risk of maternal death	United Nations Children's Fund/World Health Organization/United Nations Population Fund/ World Bank	United Nations Children's Fund/World Health Organization/United Nations Population Fund/ World Bank
	Total maternal deaths	United Nations Children's Fund/World Health Organization/United Nations Population Fund/ World Bank	United Nations Children's Fund/World Health Organization/United Nations Population Fund/ World Bank
	Maternal deaths by cause (regional)	World Health Organization	World Health Organization
Nutrition			
Anthropometry	Underweight prevalence	Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other national surveys	United Nations Children's Fund/World Health Organization
	Stunting prevalence	Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other	United Nations Children's Fund/World Health Organization
		national surveys	
	Wasting prevalence	national surveys Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other national surveys	United Nations Children's Fund/World Health Organization
Infant feeding	Wasting prevalence Exclusive breastfeeding rate (<6 months)	Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other	
Infant feeding		Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other national surveys Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other national surveys Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other	Organization
Infant feeding Low birthweight	Exclusive breastfeeding rate (<6 months)	Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other national surveys Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other national surveys Multiple Indicator Cluster Surveys,	Organization United Nations Children's Fund
	Exclusive breastfeeding rate (<6 months) Complementary feeding rate (6-9 months)	Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other national surveys Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other national surveys Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other national surveys Multiple Indicator Cluster Surveys, Demographic and Health Surveys, Demographic and Health Surveys, other	Organization United Nations Children's Fund United Nations Children's Fund
Low birthweight Micronutrient	Exclusive breastfeeding rate (<6 months) Complementary feeding rate (6-9 months) Low birthweight incidence	Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other national surveys Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other national surveys Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other national surveys Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other national surveys National Immunization Days Reporting, Demographic and Health Surveys, Multiple	Organization United Nations Children's Fund United Nations Children's Fund United Nations Children's Fund
Low birthweight Micronutrient supplementation	Exclusive breastfeeding rate (<6 months) Complementary feeding rate (6-9 months) Low birthweight incidence	Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other national surveys Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other national surveys Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other national surveys Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other national surveys Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other national surveys National Immunization Days Reporting, Demographic and Health Surveys, Multiple Indicator Cluster Survey, Routine reporting Routine reporting, Multiple Indicator Cluster Surveys, Demographic and Health Surveys	Organization United Nations Children's Fund United Nations Children's Fund United Nations Children's Fund
Low birthweight Micronutrient supplementation Child health	Exclusive breastfeeding rate (<6 months) Complementary feeding rate (6-9 months) Low birthweight incidence Vitamin A supplementation (2 doses)	Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other national surveys Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other national surveys Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other national surveys Multiple Indicator Cluster Surveys, Demographic and Health Surveys, Demographic and Health Surveys, other national surveys National Immunization Days Reporting, Demographic and Health Surveys, Multiple Indicator Cluster Survey, Routine reporting Routine reporting, Multiple Indicator Cluster	Organization United Nations Children's Fund
Low birthweight Micronutrient supplementation Child health	Exclusive breastfeeding rate (<6 months) Complementary feeding rate (6-9 months) Low birthweight incidence Vitamin A supplementation (2 doses) Measles immunization coverage Three doses of combined diphtheria/pertussis/tetanus vaccine immunization	Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other national surveys Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other national surveys Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other national surveys Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other national surveys Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other national surveys National Immunization Days Reporting, Demographic and Health Surveys, Multiple Indicator Cluster Survey, Routine reporting Routine reporting, Multiple Indicator Cluster Surveys, Demographic and Health Surveys Routine reporting, Multiple Indicator Cluster	Organization United Nations Children's Fund United Nations Children's Fund United Nations Children's Fund United Nations Children's Fund United Nations Children's Fund/World Health Organization United Nations Children's Fund/World Health
Low birthweight Micronutrient supplementation Child health	Exclusive breastfeeding rate (<6 months) Complementary feeding rate (6-9 months) Low birthweight incidence Vitamin A supplementation (2 doses) Measles immunization coverage Three doses of combined diphtheria/pertussis/tetanus vaccine immunization coverage (DPT)	Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other national surveys Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other national surveys Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other national surveys Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other national surveys Multiple Indicator Cluster Surveys, Demographic and Health Surveys, Multiple Indicator Cluster Survey, Routine reporting Routine reporting, Multiple Indicator Cluster Surveys, Demographic and Health Surveys Routine reporting, Multiple Indicator Cluster Surveys, Demographic and Health Surveys Routine reporting, Multiple Indicator Cluster Surveys, Demographic and Health Surveys	Organization United Nations Children's Fund United Nations Children's Fund United Nations Children's Fund United Nations Children's Fund United Nations Children's Fund/World Health Organization United Nations Children's Fund/World Health Organization United Nations Children's Fund/World Health Organization

Indicator		Data source	Global database
	Antimalarial treatment (children under age 5 with fever)	Multiple Indicator Cluster Surveys, Demographic and Health Surveys, Malaria Indicator Surveys	United Nations Children's Fund
Pneumonia	Careseeking for pneumonia	Multiple Indicator Cluster Surveys, Demographic and Health Surveys	United Nations Children's Fund
	Antibiotic treatment for pneumonia	Multiple Indicator Cluster Surveys, Demographic and Health Surveys	United Nations Children's Fund
Diarrhoeal diseases	Oral rehydration therapy and continued feeding	Multiple Indicator Cluster Surveys, Demographic and Health Surveys	United Nations Children's Fund
AIDS	HIV-positive pregnant women receiving antiretrovirals for prevention of mother-to- child transmission of HIV	Ministries of Health, Joint United Nations Programme on HIV/AIDS	United Nations Children's Fund
Maternal and newborn h	nealth		
Adolescent birth rate	Adolescent birth rate	Multiple Indicator Cluster Surveys, Demographic and Health Surveys, Reproductive Health Surveys, other national surveys	United Nations Population Fund/United Nations Population Division
Unmet need	Unmet need for family planning	Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other national surveys	United Nations Population Fund
Antenatal care	Antenatal care (at least one visit)	Multiple Indicator Cluster Surveys, Demographic and Health Surveys, Reproductive Health Surveys, other national surveys	United Nations Children's Fund
	Antenatal care (four or more visits)	Demographic and Health Surveys, Reproductive Health Survey, other national surveys	United Nations Children's Fund
Intermittent preventive treatment of malaria during pregnancy	Intermittent preventive treatment for pregnant women	Multiple Indicator Cluster Surveys, Demographic and Health Surveys	United Nations Children's Fund
Neonatal tetanus protection	Neonatal tetanus protection	Multiple Indicator Cluster Surveys, Demographic and Health Surveys	United Nations Children's Fund/World Health Organization
Delivery care	Skilled attendant at birth	Multiple Indicator Cluster Surveys, Demographic and Health Surveys, Reproductive Health Surveys, other national surveys	United Nations Children's Fund
Caesarean section	Caesarean section rate	Demographic and Health Surveys, Reproductive Health Survey, other national surveys	United Nations Children's Fund
Postnatal visit	Postnatal visit for mother	Demographic and Health Surveys	Special data analysis by Saving Newborn Lives
	Postnatal visit for newborn	Demographic and Health Surveys	Special data analysis by Saving Newborn Lives
Breastfeeding	Early initiation of breastfeeding	Multiple Indicator Cluster Surveys, Demographic and Health Surveys, other national surveys	United Nations Children's Fund
Contraceptive prevalence	Contraceptive prevalence rate	Multiple Indicator Cluster Surveys, Demographic and Health Surveys, Reproductive Health Surveys, other national surveys	United Nations Population Fund/United Nations Population Division
Water and sanitation			
Water	Use of improved drinking water sources	United Nations Children's Fund/World Health Organization	United Nations Children's Fund/World Health Organization
Sanitation	Use of improved sanitation facilities	United Nations Children's Fund/World Health Organization	United Nations Children's Fund/World Health Organization
Polices, systems and eq		Wald Haalth Organization (11 in 181 in	Charles data again that are health and the second
Policies	International code of marketing of breastmilk substitutes	World Health Organization/ United Nations Children's Fund	Special data compilation by World Health Organization and the United Nations Children's Fund
	New oral rehydration salts formula and zinc for management of diarrhoea	World Health Organization/ United Nations Children's Fund	Special data compilation by World Health Organization and the United Nations Children's Fund
	Community treatment of pneumonia with antibiotics	World Health Organization/ United Nations Children's Fund	Special data compilation by World Health Organization and the United Nations Children's Fund
	Integrated management of childhood illness adapted to cover newborns ages 0–1 week	World Health Organization	Special data compilation by World Health Organization
	Costed implementation plan for maternal, newborn and child health available	World Health Organization	Special data compilation by World Health Organization
	Midwives authorized to administer a core set of life-saving interventions	World Health Organization	Special data compilation by World Health Organization

Indicator		Data source	Global database
	Maternity protection in accordance with International Labour Organization Convention 183	International Labour Organization Database of Conditions of Work and Employment Laws – Maternity Protection	International Labour Organization
	Specific notification of maternal deaths	World Health Organization	Special data compilation by World Health Organization
Systems	Per capita expenditure on health	World Health Organization, <i>International Travel</i> and <i>Health Report 2010</i>	World Health Organization
	General government expenditure on health as percentage of total government expenditure	World Health Organization, <i>International Travel</i> and <i>Health Report 2010</i>	World Health Organization
	Out-of-pocket expenditure as percentage of total expenditure on health	World Health Organization, <i>International Travel</i> and <i>Health Report 2010</i>	World Health Organization
	Density of doctors, midwives and nurses per 10,000 population	World Health Organization, <i>Global Atlas of the Health Workforce</i>	World Health Organization
	Official development assistance to child health per child	Organisation for Economic Co-operation and Development/Development Assistance Committee	London School of Health and Tropical Medicine
	Official development assistance to maternal and neonatal health per live birth	Organisation for Economic Co-operation and Development/Development Assistance Committee	London School of Health and Tropical Medicine
	National availability of emergency obstetric care services	Averting Maternal Death and Disability Project and United Nations Children's Fund analysis from available emergency obstetric care surveys	Averting Maternal Death and Disability/United Nations Children's Fund
Equity	Coverage gap by wealth quintile (based on a weighted average of coverage levels of reproductive, maternal, newborn and child interventions)	Demographic and Health Surveys	Special data analysis by Federal University of Pelotas, Brazil, and the World Health Organization

Annex B Definitions of *Countdown* indicators

Intervention	Indicator definition	Numerator	Denominator
Nutrition			
Exclusive breastfeeding (<6 months)	Percentage of infants ages 0–5 months who are exclusively breastfed	Number of infants ages 0-5 months who are exclusively breastfed	Total number of infants ages 0—5 months surveyed
Complementary feeding (6–9 months)	Percentage of infants ages 6–9 months who are breastfed and receive complementary food	Number of infants ages 6–9 months who are breastfed and receive complementary food	
Vitamin A supplementation	Percentage of children ages 6–59 months who received two doses of vitamin A during the calendar year	Number of children ages 6–59 months who received two doses of vitamin A during the calendar year	Total number of children ages 6–59 months
Child health			
Measles immunization coverage	Percentage of infants immunized with measles containing vaccine	Number of children ages 12–23 months who are immunized against measles	Total number of children ages 12–23 months surveyed
Three doses of combined diphtheria/ pertussis/tetanus (DPT) vaccine immunization coverage	Percentage of infants who received three doses of diphtheria/pertussis/tetanus vaccine	Number of children ages 12–23 months receiving three doses of diphtheria/pertussis/tetanus vaccine	Total number of children ages 12–23 months surveyed
Three doses of <i>Haemophilus influenzae</i> type B immunization coverage	Percentage of infants who received three doses of <i>Haemophilus influenzae</i> type B vaccine	Number of children ages 12–23 months receiving three doses of <i>Haemophilus influenzae</i> type B vaccine	Total number of children ages 12–23 months surveyed
Oral rehydration therapy and continued feeding	Percentage of children ages 0–59 months with diarrhoea receiving oral rehydration therapy and continued feeding	Number of children ages 0–59 months with diarrhoea in the two weeks prior to the survey receiving oral rehydration therapy (oral rehydration solution and/or recommended homemade fluids or increased fluids) and continued feeding	Total number of children ages 0–59 months with diarrhoea in the two weeks prior to the survey
Insecticide-treated net use	Percentage of children ages 0-59 months sleeping under an insecticide-treated mosquito net	Number of children ages 0–59 months sleeping under an insecticide-treated mosquito net the night before the survey	Total number of children ages 0–59 months surveyed
Antimalarial treatment	Percentage of children ages 0—59 months with fever receiving any appropriate antimalarial drugs	Number of children ages 0–59 months reported to have fever in the two weeks prior to the survey who were treated with any appropriate antimalarial	Total number of children ages 0–59 months reported to have fever in the two weeks prior to the survey
Prevention of mother-to-child transmission of HIV	Percentage of HIV-infected pregnant women who received antiretrovirals to reduce the risk of mother-to-child transmission	Number of HIV-infected pregnant women who received antiretrovirals in the last 12 months to reduce mother-to-child transmission	Estimated unrounded number of HIV-infected pregnant women ^a
Careseeking for pneumonia	Percentage of children ages 0–59 months with suspected pneumonia taken to an appropriate health provider	Number of children ages 0–59 months with suspected pneumonia in the two weeks prior to the survey who were taken to an appropriate health provider	Total number of children ages 0–59 months with suspected pneumonia in the two weeks prior to the survey
Antibiotic treatment for pneumonia	Percentage of children ages 0–59 months with suspected pneumonia receiving antibiotics	Number of children ages 0–59 months with suspected pneumonia in the two weeks prior to the survey receiving antibiotics	Total number of children ages 0–59 months with suspected pneumonia in the two weeks prior to the survey
Maternal and newborn health			
Contraceptive prevalence	Percentage of women currently married or in union ages 15–49 that are using (or whose partner is using) a contraceptive method (either modern or traditional)	Number of women currently married or in union ages 15–49 years that are using (or whose partner is using) a contraceptive method (either modern or traditional)	Total number of women ages 15–49 that are currently married or in union
Adolescent birth rate	Number of births to adolescent women per 1,000 adolescent women ^b	Number of live births to adolescent women ages 15–19	1,000 adolescent women ages 15–19
Unmet need for family planning	Percentage of women who are currently married or in union that have an unmet need for contraception	Number of women who are currently married or in union that are fecund and want to space their births or limit the number of children they have but that are not currently using contraception	Total number of women who are currently married or in union
Antenatal care (at least one visit)	Percentage of women attended at least once during pregnancy by skilled health personnel for reasons related to the pregnancy	Number of women attended at least once during pregnancy by skilled health personnel (doctor, nurse, midwife or auxiliary midwife) for reasons related to the pregnancy in the X years prior to the survey	Total number of women who had a live birth occurring in the same period
Antenatal care (four or more visits)	Percent of women attended at least four times during pregnancy by any provider (skilled or unskilled) for reasons related to the pregnancy	to the pregnancy in the X years prior to the survey	Total number of women who had a live birth occurring in the same period
Neonatal tetanus protection	Percentage of newborns protected against tetanus	Number of mothers with a live birth in the year prior to the survey who received two doses of tetanus toxoid vaccine within the appropriate interval prior to the infant's birth	

Intervention	Indicator definition	Numerator	Denominator
Intermittent preventive treatment for malaria during pregnancy	Percentage of women who received intermittent preventive treatment for malaria during their last pregnancy	Number of women at risk for malaria who received two or more doses of a sulfadoxine-pyrimethamine (Fansidar TM) to prevent malaria during their last pregnancy that led to a live birth	Total number of women surveyed who delivered a live newborn within the last two years
Skilled attendant at birth	Percentage of live births attended by skilled health personnel	Number of live births to women ages 15–49 years in the X years prior to the survey attended during delivery by skilled health personnel (doctor, nurse, midwife or auxiliary midwife)	Total number of live births to women ages 15–49 years in the X years prior to the survey ^c
Caesarean section rate	Percentage of live births delivered by Caesarean section	Number of live births to women ages 15–49 years in the X years prior to the survey delivered by caesarean section	Total number of live births to women ages 15–49 years in the X years prior to the survey
Early initiation of breastfeeding	Percentage of newborns put to the breast within one hour of birth	Number of women with a live birth in the X years prior to the survey who put the newborn infant to the breast within 1 hour of birth	Total number of women with a live birth in the X years prior to the survey ^d
Postnatal care for mothers ^e	Percentage of mothers who received postnatal care visit within two days of childbirth	Number of women who received a postnatal care visit within two days of childbirth (regardless of place of delivery)	Total number of women ages 15–49 years with a last live birth in the x years prior to the survey (regardless of place of delivery)
Postnatal care for babies who were born at home	Percentage of babies born outside a facility who received a postnatal care visit within two days of birth	Number of babies born outside of a health facility who received a postnatal care visit within two days of birth ^f	Total number of last-born babies born outside of a health facility in the X years prior to the survey ^g
Water and sanitation			
Use of improved drinking water sources	Percentage of the population using improved drinking water sources	Number of household members living in households using improved drinking water sources (including household connections, public standpipe, borehole, protected dug well, protected spring, rainwater collection)	Total number of household members in households surveyed
Use of improved sanitation facilities	Percentage of the population using improved sanitation facilities	Number of household members using improved sanitation facilities (including connection to a public sewer, connection to a septic system, pour-flush latrine, simple pit latrine, or a ventilated improved pit latrine)	Total number of household members in households surveyed

Notes

- a. More details on the HIV estimates methodology can be found at www.unaids.org.
- b. Also referred to as the age-specific fertility rate for women ages 15–19.
- $\ensuremath{\text{c.}}$ The reference period may differ between surveys.
- d. The reference period may differ between surveys.
- e. As used for postnatal care in figure 4.
- f. Information on postnatal care for babies who were born in health facilities has historically not been collected by Demographic and Health Surveys.
- g. This denominator differs from the all births denominator used for the indicator for postnatal care for mother. Therefore, the coverage for mother and newborn cannot be compared.

Annex C Definitions of policy and health systems indicators

Indicator	Indicator definition	Criteria for ranking
Midwives authorized to administer a core set of life-	National policy adopted authorizing midwives to administer the following:	Yes: midwives authorized for all tasks
saving interventions	parental antibioticsparental oxytocics	Partial: midwives authorized for some tasks
	parental anticonvulsants manual removal of placenta removal of retained products of conception assisted vaginal delivery newborn resuscitation	No: midwives not authorized for any of these tasks
Specific notification of maternal deaths	National policy adopted requiring health professionals to notify any maternal death	Yes: national policy adopted and implemented
deaths	professionals to notify any material accuti	Partial: national policy adopted but no systematic implementation
late metal management of	Markey distance to decrease a fight disease	No: no national policy
Integrated management of childhood illness adapted to cover newborns ages 0–1 week	National integrated management of childhood illness guidelines adapted to cover major conditions affecting newborn survival in the first week of life	Yes: National integrated management of childhood illness guidelines adapted and in line with World Health Organization generic guidelines 2006
		Partial: National integrated management of childhood illness guidelines adapted but not fully in line with World Health Organization generic guidelines 2006
		No: National integrated management of childhood illness guidelines not adapted
New oral rehydration salts formula and zinc for	National policy guidelines adopted on management of diarrhoea with low osmolarity oral rehydration	Yes: low osmolarity oral rehydration salts and zinc supplements in national policy
management of diarrhoea	salts and zinc supplements	Partial: low osmolarity oral rehydration salts or zinc supplements in national policy
		No: low osmolarity oral rehydration salts and zinc supplements not promoted in national policy
Community treatment of pneumonia with antibiotics	National policy adopted authorizing community health workers to identify and manage pneumonia	Yes: community health workers authorized to give antibiotics for pneumonia
	with antibiotics	Partial: no national policy but some implementation of community-based management of pneumonia
		No: no national policy and no implementation
Maternity protection in accordance with International Labour Organization Convention 183	ection in International Labour Organization Convention 183 th International ratified by the country or national legislation in	Yes: International Labour Organization Convention 183 ratified (maternity leave of at least 14 weeks with cash benefits of previous earnings paid by social security or public funds)
	maternity leave and level and source of funding of maternity benefits—namely 14 weeks paid at 66% of previous earnings by social security or general revenue)	Partial: International Labour Organization Convention 183 not ratified but previous maternity convention ratified (maternity leave of at least 12 weeks with cash benefit paid by social security or public funds)
		No: no ratification of any maternity protection convention
International Code of Marketing of Breastmilk Substitutes	National policy adopted on all provisions stipulated in the International Code of Marketing of Breastmilk Substitutes	Yes: all provisions of the International Code of Marketing of Breastmilk Substitutes adopted in legislation
		Partial: voluntary agreements or some provisions of the International Code of Marketing of Breastmilk Substitutes adopted in legislation
		No: no legislation and no voluntary agreements adopted in relation to the International Code of Marketing of Breastmilk Substitutes
Costed implementation plan for maternal, newborn and child health	National plan or plans for scaling up maternal, newborn and child health interventions available and costed	Yes: costed plan or plans to scale up maternal, newborn and child health interventions available at national level
		Partial: costed plan available for either maternal and newborn health or child health
		No: no costed implementation plan for maternal, newborn and child health available
Per capita expenditure on health (at international dollar rate)	Data on per capita health expenditure in international dollars for 2004 in this edition of <i>Countdown</i> are not comparable with data in previous editions because the estimates in this edition are based on the new purchasing power parity series from the World Bank's 2005 International Comparison Program. Data for 2004 and 2006 in this edition are fully comparable.	
Per capita expenditure on health as percentage of total government expenditure		Numerical

Indicator	Indicator definition	Criteria for ranking
Out-of-pocket expenditure as percentage of total expenditure on health		Numerical
Density of health workers per 10,000 population	Total number of physicians, nurses and midwives relative to the overall population	Numerical minimum standard: 23 health workers per 10,000 people needed to deliver basic maternal and child health services
Availability of emergency obstetric care services as percentage of recommended minimum	care facilities per 500,000 people, including one comprehensive and four basic emergency obstetric care facilities. (The breakdown of comprehensive and basic by population and geographic area is available	Availability is expressed as a percentage of the minimum acceptable number of emergency obstetric care facilities
		The minimum acceptable number of emergency obstetric care facilities (comprehensive and basic) is calculated by dividing the population by 500,000 and multiplying by 5
		The percentage of recommended minimum number of emergency obstetric care facilities is calculated by dividing the number of functioning emergency obstetric care facilities by the recommended number and multiplying by 100. To qualify as a fully functioning basic or comprehensive emergency obstetric care facility, a facility must provide a standard set of signal functions

Annex D

Technical notes on measuring equity in reproductive, maternal, newborn and child health through the coverage gap index

Coverage indicators

The measure of equity constructed for the *Countdown* report is the coverage gap index. For guidance on interpreting the coverage gap graphs in the country profiles, please see the section on explanation and interpretation of coverage gap below. The coverage gap index combines information on four intervention areas across the continuum of care: family planning, maternal and newborn care, immunization and treatment of sick children. Data from Demographic and Health Surveys and Multiple Indicator Cluster Surveys on eight coverage indicators in these four intervention areas was used to construct the coverage gap index (table D1).

Calculation of the coverage gap index

The coverage gap index was calculated using the formula:

100% - ([ORT + ARI] / 2 + FPS + [SBA + ANC] / 2 + [MSL + 2*DPT3 + BCG] / 4) / 4

Each of the four intervention areas is given equal weight.

If need data on need for family planning satisfied were not available, the contraceptive prevalence

rate among married women ages 15–49 was used to estimate the need satisfied according to the following formula: FPS = CPR * 1.07 + 27. This formula was derived from analysis of more than 100 Demographic and Health Surveys with data on both unmet need and contraceptive prevalence rate.

Wealth index

The coverage gap index was calculated for the total sample for each country and data point. To measure equity, the total sample must be divided into groups by socioeconomic status. Demographic and Health Surveys and Multiple Indicator Cluster Surveys do not collect information on income and expenditure, which could be used to do this. But the surveys do collect information on asset ownership and availability of basic household services. For the purposes of analyzing socioeconomic inequalities in health, it has been shown that using such variables to develop an index of socioeconomic status leads to similar results as using income or expenditure data.⁴⁰

For coverage of health interventions in the Demographic and Health Surveys, data from an analysis by Gwatkin and others (2007) were used. They took information on household assets and access to basic household services from Demographic and Health Surveys to construct a wealth index. The index was used to rank households and then divide the household population into quintiles. Results from recent

	LE D1 verage gap index indicator de	efinitions
Number	Indicator	Definition
1a.	Need for family planning satisfied (FPS)	Percentage of currently married women who are fecund and say that they do not want any more children or that they want to wait two or more years before having another child or who had a unwanted pregnancy in the previous six months that are using contraception (any method)
1b.	Contraceptive prevalence rate (CPR)	Percentage of women ages 15–49 currently married or in union who are using (or whose partner is using) a modern contraceptive method
2.	Antenatal care (ANC)	Percentage of women attended at least once during pregnancy by skilled health personnel for reasons relate to the pregnancy in the three years prior to the survey
3.	Skilled birth attendance (SBA)	Percentage of live births in the three years prior to the survey attended by skilled health personnel (doctor, nurse, midwife or auxiliary midwife)
4.	Measles vaccination (MSL)	Percentage of children ages 12–23 months who are immunized against measles
5.	Diphtheria, pertussis and tetanus vaccination (three doses of combined DPT vaccine)	Percentage of children ages 12–23 months who received three doses of DPT vaccine
6.	BCG vaccination (BCG)	Percentage of children age 1–23 months currently vaccinated against BCG
7.	Oral rehydration therapy (ORT)	Percentage of children under age 5 with diarrhoea in the last two weeks who received oral rehydration therap (oral rehydration salts packets, recommended home solution or increased fluids) and continued feeding
8.	Treatment of acute respiratory infection (ARI)	Percentage of children ages 0–59 months with suspected pneumonia (cough and dyspnoea) who sought care from a health provider

Demographic and Health Surveys results were also included. For Multiple Indicator Cluster Surveys, data from the United Nations Children's Fund's Multiple Indicator Cluster Surveys website (http://childinfo.org) were used for those countries as well as data points for which a wealth index had been constructed.⁴¹

Explanation and interpretation of coverage gap graph

The *x*-axis shows the wealth quintiles; from the poorest quintile to the best-off quintile. The *y*-axis shows the mean coverage of the selected interventions (in green, bottom part of the area chart) and the coverage gap (in red, top part of the area chart). The coverage gap represents how much the mean coverage is below 100%.

The difference between the poorest and richest quintiles and the shape of the line show the patterns of inequality within a country. First, the greater the inequality between the poorest and richest quintiles, the steeper the upward slope of the mean coverage line. With a few exceptions, the green area increases moving from the poorest quintile to the best-off quintile in the country profiles, showing higher coverage (and therefore a smaller gap) among the better off.

The shape of the area is equally important.⁴² The way the lines are curved can illustrate where

inequities are concentrated. There are three main patterns. First, bottom inequity occurs when the poorest lag behind. Second, top inequity occurs when the richest do substantially better than the other quintiles. The intermediate pattern is more or less linear. Coverage increases by a similar fraction as one goes from the poorest to the richest quintile.

The shape of the coverage gap line can inform policies to address inequities. Many country graphs have relatively straight upward-sloping lines from the poorest to the best-off quintile, which would suggest that efforts should be made to increase the overall coverage of interventions, but with special attention paid to the poor. A top inequity pattern, as illustrated in the Nigeria and Uganda country profiles, with a relatively small coverage gap among the best-off quintile, suggests that inequities would be reduced by raising the overall population coverage of interventions.

An upward slope from the poorest quintile to the second-poorest quintile and then a more or less straight line (or at least less steep) to the best-off quintile would be an example of bottom inequity, as shown in the South Africa country profile. Such a pattern indicates that inequities are concentrated among the poorest and that the most appropriate policy response would be to target that particular group.

Annex E Countdown priority countries considered to be malaria endemic

Table E1 indicates which *Countdown* priority countries are malaria endemic—defined as having a documented risk of *Plasmodium falciparum* transmission nationwide and throughout the year—and which countries have subnational risk, mostly *p. vivax*, no risk or very limited risk.

TABLE E1	
Plasmodium falciparum transmission risk in Count	down countries

Malaria-endemic countries (n = 45)		Countries with subnational risk of Plasmodium falciparum transmission (n = 14)	Countries with mostly p. vivax, no Plasmodium falciparum or very limited risk (n = 9)	
Afghanistan Angola Bangladesh Benin Botswana Burkina Faso Burundi Cambodia ^a Cameroon Central African Republic Chad Congo Congo, Dem. Rep. of the Côte d'Ivoire Djibouti Equatorial Guinea Eritrea Ethiopia ^a Gabon Gambia Ghana Guinea Guinea Guinea	Kenya ^a Lao PDR Liberia Madagascar Malawi Mali Mozambique Myanmar ^a Niger Nigeria Pakistan ^a Papua New Guinea ^a Rwanda Senegal ^a Sierra Leone Somalia Sudan ^a Tanzania, U. Rep. Togo Uganda Zambia Zimbabwe	Bolivia Brazil China Haiti India Indonesia Mauritania Nepal Peru Philippines South Africa Swaziland Tajikistan Yemen	Azerbaijan Egypt Guatemala Iraq Korea, Dem. Rep. Lesotho Mexico Morocco Turkmenistan	

a. Country has lower risk of *Plasmodium falciparum* transmission in identifiable areas (such as certain urban centres) but the highest prevention strategy is still recommended nationwide.

Source: WHO 2010.

Annex F Details on estimates from the Inter-agency Group for Child Mortality Estimation used in the *Countdown* report

Child mortality estimates

The child mortality estimates in this report (infant mortality rate, under-five mortality rate and under-five deaths) are based on the work of the Inter-agency Group for Child Mortality Estimation, which includes the United Nations Children's Fund (UNICEF), the World Health Organization (WHO), the United Nations Population Division and the World Bank. The estimates are the official UN estimates for measuring progress towards Millennium Development Goal 4-reducing child mortality. The Inter-agency Group for Child Mortality Estimation compiles data available from all possible nationally representative sources for a country, including household surveys, censuses, vital registration and then like, and uses a model to fit a regression line to the data to produce the mortality estimates. Estimates are updated every year after a detailed review of all newly available data points. The review occasionally results in adjustments to previously reported estimates. The full time series for all countries is published at www.childinfo.org and www.childmortality.org.

Maternal mortality estimates

Maternal mortality estimates for 2005 are based on the work of an interagency group comprising the WHO, UNICEF, the United Nations Population Fund and the World Bank. The model-based estimates use a dual approach to take into account the frequent underreporting and misclassification of maternal deaths. For more information, see WHO, UNICEF, UNFPA and World Bank (2007).

Immunization

The immunization data published in this report are based on the work of the WHO and UNICEF. The estimates should not be confused with other sources of information, such as Demographic and Health Surveys, Multiple Indicator Cluster Surveys, or Ministries of Health. The WHO and UNICEF derive national estimates of immunization coverage each year and send each country a draft report for review. Based on comments and new evidence from the country reviews, final reports are published in August with data for the preceding year. All new evidence, such as final survey reports received after publication, are taken into consideration for the following year's estimates. For each country's final report for 2008 as well as data sources and a description of trends, see www.childinfo.org.

Water and Sanitation

The drinking water and sanitation coverage estimates are produced by the WHO-UNICEF Joint Monitoring Programme for Water Supply and Sanitation. The estimates are the official UN estimates for measuring progress towards the Millennium Development Goal targets for drinking water and sanitation. They use a standard classification of what constitutes coverage. The Joint Monitoring Programme does not report the findings of the latest nationally representative household survey or census. Instead, it estimates coverage using a linear regression line that is based on coverage data from all available household sample surveys and census. For specific country data, see www.childinfo.org and www. wssinfo.org.

Annex G Calculating the financing gap for maternal, newborn and child health

To project likely trends of financial flows to maternal, newborn and child health over 2008–2015, a spreadsheet model was developed covering two scenarios: a business as usual scenario based on past experience and a public commitments scenario based on national government and bilateral donor commitments (see table G1 for the main assumptions of the model).

To estimate the financing gap, forecasted expenditure was compared with costs of scaling

up maternal, newborn and child interventions. The costs used were estimated by the Millennium Development Goals 4 and 5 costing and impact estimate group for the Global Campaign for the Health Millennium Development Goals (2008).

Data were converted from nominal values in national currency units into real data with a 2007 base year using country-specific consumer price index growth rates and then converted into U.S. dollars using International Monetary Fund exchange rates.

The complete aid activities database for 2007 was analyzed and coded according to the methodology used in Powell-Jackson and others

TABLE G1

Assumptions of the two scenarios in the model for projecting financial flows to maternal, newborn and child health

Variable	Scenario 1: business as usual	Scenario 2: public commitments	Source
Panel A: General government health ex	xpenditure		
GDP forecast	Country-specific annual growth rates		International Monetary Fund <i>World Economic Outlook</i>
Ratio of general government expenditure to GDP	Average 2000–2007		World Health Organization and International Monetary Fund historic dat
Ratio of general government expenditure for health to general government expenditure	Average 2000–2007	Linear increase to 15% for African countries Linear increase to 12% for non-African countries	World Health Organization historic data Abuja Declaration
Distribution of external health expenditure among public and private financing agents	70% government financing agents 30% private financing agents		Mean of estimates from national health accounts reports
Panel B: Private health expenditure	co /o pate intaileing agents		
Private health expenditure	Increases in line with real GDP growth fro	om 2007 baseline value	Best estimate
Distribution of external health expenditure among public and private financing agents	70% government financing agents 30% private financing agents		Mean of estimates from national health accounts reports
Panel C: external health expenditure	50% private infancing agents		
Annual real GDP growth in donor countries	2% a year		Organisation for Economic Co-operation and Development/Development Assistance Committee projections of net official development assistance
Ratio of official development assistance to GDP in donor countries	2008–2015 remains constant at 2007 ratio	Linear increase to 0.7% in 2015, or the ratio committed to in 2010, whichever is higher. Exceptions are Japan and the United States: linear increase to 0.3% in 2015	Organisation for Economic Co-operation and Development/Development Assistance Committee 2007 data Organisation for Economic Co-operation and Development/Development Assistance Committee projections of net official development assistance
Distribution of official development assistance to health and across study countries	2008–2015 remains constant at 2007 app	Organisation for Economic Co-operation and Development Creditor Reporting System 2007 data	
Panel D: maternal, newborn and child	health expenditure		
Ratio of maternal, newborn and child health expenditure to health expenditure	Government expenditure: 25%		National health accounts reports
	Private expenditure: 25%		
	External expenditure: 25%		

(2006) and Greco and others (2010). This is a preliminary analysis; a final analysis of the 2007 and 2008 data will be completed by December 2010.

All 22 donor countries and the EU members of the Organisation for Economic Co-operation and Development's Development Assistance Committee were included. In addition, the World Bank, UNICEF, the Joint United Nations Programme on HIV/AIDS, the Global Alliance for Vaccines and Immunisation and the Global Fund to Fight AIDS, Tuberculosis and Malaria were included as multilateral development organizations and global health funds. After two years of nonreporting, the United Nations Population Fund

reported data for 2007, so its disbursements are also included.

For all but one donor, the analysis used data from the Organisation for Economic Co-operation and Development Creditor Reporting System database (www.oecd.org/dac/stats/idsonline). The Global Alliance for Vaccines and Immunisation provided disbursement data, which were also included in the analysis. Disbursements from the Global Fund to fight AIDS, Tuberculosis and Malaria were available in the Creditor Reporting System database, but data were cross-checked against the data on the Global Fund website and found to be consistent. The World Bank did not report disbursements, so its commitment data were used instead.

Notes



- 1. You and others 2009.
- 2. Jones and others 2003.
- 3. Black and others forthcoming.
- 4. Black and others forthcoming.
- 5. Gwatkin and others 2007.
- 6. Rutstein 2005.
- 7. WHO 1985.
- 8. Wall and others 2009.
- 9. Mwansa-Kambafwile and others 2010.
- 10. Lawn and others 2010.
- 11. WHO and UNICEF 2009.
- 12. UNICEF and WHO 2009.
- 13. UNICEF, Roll Back Malaria, and The Global Fund to Fight AIDS, Tuberculosis and Malaria 2009.
- 14. WHO 2006.
- 15. WHO 2009.
- 16. Phillips, Bawah, and Binka 2006; Wakabi 2008; Douthwaite and Ward 2005.
- 17. Dovlo 2004.
- 18. Pereira and others 2007; Vaz and others 1999.
- 19. WHO 2010.
- 20. Xu and others 2005.
- 21. WHO 2005b.

- 22. WHO 2005a.
- 23. Gonzalez-Pier and others 2007.
- 24. Parry and Weiyuan 2008.
- 25. Kagumire 2009.
- 26. Franco and others 2008.
- 27. Nabyonga and others 2005.
- 28. SUZY Project, ICDDR,B 2009 (personal communication).
- 29. Shankar and others 2008.
- 30. AbouZahr and others 2007.
- 31. WHO Regional Office for South-East Asia 2009.
- 32. South Africa Every Death Counts Writing Group 2008.
- 33. WHO and UNICEF 2004a, 2004, 2009.
- 34. Dawson and others 2008.
- 35. Koblinsky and others 1999, 2006.
- 36. Boerma and others 2008.
- 37. UNICEF 2007.
- 38. UNICEF 2007.
- 39. Bishai and others 2002.
- 40. Wagstaff and Watanabe 2003.
- 41. For more information on the calculation of the wealth index from DHS and MICS data, please refer to Rutstein and Johnson (2004).
- 42. Victora and others 2005.

References



- AbouZahr, C., and others. 2007. "The Way Forward." *Lancet* 370 (9601): 1791–1799.
- Alliance for Health Policy and System Research. 2010. Systems Thinking for Health System Strengthening. Geneva: World Health Organization.
- Bang, A.T. and others. 1999. "Effect of Home-Based Neonatal Care and Management of Sepsis on Neonatal Mortality: Field Trial in Rural India." *Lancet* 354 (9194): 1955–1961.
- Baqui, A.H., and others. 2009. "Effectiveness of Home-Based Management of Newborn Infections by Community Health Workers in Rural Bangladesh." *Pediatric Infectious Disease Journal* 28 (4): 304–310.
- Barros, F.C., and others. 2010. "Recent Trends in Maternal, Newborn and Child Health in Brazil: A Case-Study in Progress towards MDGs 4 and 5." Unpublished draft. Pan American Health Organization, Washington, D.C.
- Basinga, P., and others. 2010. "Paying Primary Health Care Centers for Performance in Rwanda." Policy Research Working Paper 5190. World Bank, Washington, D.C.
- Bishai, D., and others. 2002. "The Role of Public Health Programmes in Reducing Socioeconomic Inequities in Childhood Immunization Coverage." *Health Policy Plan* 17 (4): 412–419.
- Black, R.E., and others. 2008. "Maternal and Child Undernutrition: Global and Regional Exposures and Health Consequences." *Lancet* 371 (9608): 243–260.
- ——. Forthcoming. "Global, Regional and National Causes of Child Deaths, 2008." *Lancet*.
- Boerma, J.T., and others. 2008. "Mind the Gap: Equity and Trends in Coverage of Maternal,

- Newborn, and Child Health Services in 54 Countdown Countries." *Lancet* 371 (9620): 1259–1267.
- Dawson P., and others. 2008. "From Research to National Expansion: 20 Years' Experience of Community-Based Management of Childhood Pneumonia in Nepal." Bulletin of the World Health Organization 86 (5): 339–343.
- Devadasan, N., and others. 2008. "A Conditional Cash Assistance Programme for Promoting Institutional Deliveries among the Poor in India: Process Evaluation Results." In F. Richard, S. Witter, and V. De Brouwere, eds., Reducing Financial Barriers to Obstetric Care in Low-Income Countries. Antwerp, Belgium: ITGPress.
- Douthwaite, M., and P. Ward. 2005. "Increasing Contraceptive Use in Rural Pakistan: An Evaluation of the Lady Health Worker Programme." *Health Policy Plan* 20 (2): 117–123.
- Dovlo, D. 2004. "Using Mid-Level Cadres as Substitutes for Internationally Mobile Health Professionals in Africa: A Desk Review." Human Resources for Health 2 (7).
- Filippi V., and others. 2006. "Maternal Health in Poor Countries: The Broader Context and a Call for Action." Lancet 368 (9546): 1535–1541.
- Franco, L.M., and others. 2008. "Effects of Mutual Health Organizations on Use of Priority Health-Care Services in Urban and Rural Mali: A Case-Control Study." Bulletin of the World Health Organization 86 (11): 830–838.
- Global Campaign for the Health Millennium
 Development Goals. 2008. First Year Report:
 2008. New York: Global Campaign for the Health
 Millennium Development Goals.
- Gonzalez-Pier, E., and others. 2007. "Priority Setting for Health Interventions in Mexico's System of

- Social Protection in Health." Salud Pública de México 49 (Suppl. 1): S37–S52.
- Greco G., T. Powell-Jackson, J. Borghi, and A. Mills. 2008. "Countdown to 2015: Assessment of Donor Assistance to Maternal, Newborn, and Child Health between 2003 and 2006." *Lancet* 371 (9620):1268–1275.
- Gwatkin, D.R., and others. 2007. "Socio-economic Differences in Health, Nutrition and Population within Developing Countries: An Overview." World Bank, Washington, D.C.
- Hogan, M.C., and others. 2010. "Maternal Mortality for 181 Countries, 1980–2008: A Systematic Analysis of Progress toward Millennium Development Goal 5." *Lancet* 375 (9726): 1609–1623.
- India, Ministry of Health and Family Welfare. 2005. *Janani Suraksha Yojana: Guidelines for Implementation*. New Delhi: Government of India.
- Jones, G., and others. 2002. "How Many Child Deaths Can We Prevent This Year?" *Lancet* 362 (9377): 65–71.
- Kagumire, R. 2009. "Public Health Insurance in Uganda Still Only a Dream." Canadian Medical Association Journal 180 (3): 281.
- Koblinsky, M.A. and others. 1999. "Organizing Delivery Care: What Works for Safe Motherhood." Bulletin of the World Health Organization 77 (5): 399–406.
- Koblinsky, M.A. and others. 2006. "Going to Scale with Professional Skilled Care." *Lancet* 368 (9544): 1377–1386.
- Lawn, J.E., K. Shibuya, and C. Stein. 2005. "No Cry at Birth: Global Estimates of Intrapartum Stillbirths and Intrapartum-Related Neonatal Deaths." Bulletin of the World Health Organization 83 (6): 409–417.
- Lawn, J.E., and others. 2009. "Two Million Intrapartum-Related Stillbirths and Neonatal Deaths: Where, Why, and What Can Be Done?" International Journal of Gynecology and Obstetrics 107 (Suppl. 1): S5–S19.
- Lawn, J.E., and others. 2010. "'Kangaroo Mother Care' to Prevent Neonatal Deaths due to Preterm Birth Complications." *International Journal of Epidemiology* 39 (Suppl. 1): i144–i154.

- Lee, A.C., and others. 2009. "Linking Families and Facilities for Care at Birth: What Works to Avert Intrapartum-Related Deaths?" International Journal of Gynecology and Obstetrics 107 (Suppl.): S65–S68.
- Li, X.F., J.A. Fortney, M. Kotelchuck, and L.H. Glover. 1996. "The Postpartum Period: The Key to Maternal Mortality." *International Journal of Gynecology and Obstetrics* 54 (1): 1–10.
- Mavalankar D., and others. 2009. "Saving Mothers and Newborns through an Innovative Partnership with Private Sector Obstetricians: Chiranjeevi Scheme of Gujarat, India." *International Journal of Gynecology and Obstetrics* 107 (Suppl.): 271–276.
- Monteiro, C.A., and others. 2010. "Narrowing Socioeconomic Inequality in Child Stunting:The Brazilian Experience, 1974–2007." Bulletin of the World Heatlh Organization 88 (4): 305–311.
- Mwansa-Kambafwile, J., S. Cousens, T. Hansen, and J.E. Lawn. 2010. "Antenatal Steroids in Preterm Labour for the Prevention of Neonatal Deaths due to Complications of Preterm Birth." *International Journal of Epidemiology* 39 (Suppl. 1): i122–133.
- Nabyonga, J., and others. 2005. "Abolition of Cost-Sharing Is Pro-Poor: Evidence from Uganda." Health Policy Plan 20 (2): 100–108.
- Parry, J., and C. Weiyuan. 2008. "Making Health Care Affordable in China." *Bulletin of the World Health Organization* 86 (11): 821–823.
- Pereira, C, and others. 2007. "Meeting the Need for Emergency Obstetric Care in Mozambique: Work Performance and Histories of Medical Doctors and Assistant Medical Officers Trained for Surgery." BJOG 114 (12): 1530–1533.
- Phillips, J.F., A.A. Bawah, and F.N. Binka. 2006. "Accelerating Reproductive and Child Health Programme Impact with Community-Based Services: The Navrongo Experiment in Ghana." Bulletin of the World Health Organization 84 (12): 949–955.
- Pitt, C., G. Greco, T. Powell-Jackson, and A. Mills. 2010. "Countdown to 2015: Tracking Donor Flows to MNCH in 2007." Draft report. London School of Hygiene and Tropical Medicine, Health Economics and Financing Programme, London.
- Powell-Jackson, T., and others 2006. "Countdown to 2015: Tracking Donor Assistance to Maternal,

- Newborn and Child Health." *Lancet* 368 (9541): 1077–1087.
- Rutstein, S.O. 2005. "Effects of Preceding Birth Intervals on Neonatal, Infant, and Under Five Years Mortality and Nutritional Status in Developing Countries: Evidence from the Demographic and Health Surveys." Journal of Gynecology and Obstetrics 89 (Suppl. 1): S7–S24.
- Rutstein, S.O., and K. Johnson. 2004. *The DHS Wealth Index*. DHS Comparative Report 6. Calverton, Md.: ORC Macro.
- Shankar, A. and others. 2008. "Delivery of MDG 5 by Active Management with Data." *Lancet* 371 (9620): 1223–1224.
- South Africa Every Death Counts Writing Group. 2008. "Every Death Counts: Use of Mortality Audit Data for Decision Making to Save the Lives of Mothers, Babies, and Children in South Africa." *Lancet* 371 (9620): 1294–1304.
- Stanton, C. 2006. "Stillbirth Rates: Delivering Estimates in 190 Countries." *Lancet* 367 (9521): 1487–1494.
- Stewart, C.P., K.G. Dewey, and P. Ashorn. 2010. "The Undernutrition Epidemic: An Urgent Health Priority. (Comment)." *Lancet* 375 (971): 282.
- Theodoratou, E., and others. 2010. "The Effect of Case Management on Childhood Pneumonia Mortality in Developing Countries." *International Journal of Epidemiology* 39 (Suppl. 1): 155–171.
- UNICEF (United Nations Children's Fund). 2007.

 Progress for Children: A World Fit for Children
 Statistic Review. New York: United Nations
 Children's Fund.
- ——. 2009a. State of the World's Children: Special Edition. New York: United Nations Children's Fund.
- . 2009b. Tracking Progress on Child and Maternal Nutrition: A Survival and Development Priority. New York: United Nations Children's Fund.
- UNICEF (United Nations Children's Fund) and WHO (World Health Organization). 2009. *Diarrhoea:* Why Children Are Still Dying and What Can Be Done. New York: United Nations Children's Fund.
- UNICEF (United Nations Children's Fund), Roll Back Malaria, and The Global Fund to Fight AIDS, Tuberculosis and Malaria. 2009. *Malaria and*

- Children: Progress in Intervention Coverage. Summary update 2009. New York: United Nations Children's Fund.
- UNICEF (United Nations Children's Fund), UNAIDS (Joint United Nations Programme on HIV/AIDS), WHO (World Health Organization), and UNFPA (United Nations Population Fund). 2009. Children and AIDS: Fourth Stocktaking Report 2009. New York and Geneva: United Nations Children's Fund, Joint United Nations Programme on HIV/AIDS, World Health Organization, and United Nations Population Fund.
- Vaz, F., and others. 1999. "Training Medical Assistants for Surgery: Policy and Practice." *Bulletin of the World Health Organization* 77 (8): 688–691.
- Victora, C.G., B. Fenn, J. Bryce, and B.R. Kirkwood. 2005. "Co-coverage of Preventive Interventions and Implications for Child Survival Strategies: Evidence from National Surveys." *Lancet* 366 (9495):1460–1466.
- Victora, C.G., and others. 2008. "Maternal and Child Undernutrition: Consequences for Adult Health and Human Capital." *Lancet* 371 (9609): 340–357.
- Wagstaff, A., and N. Watanabe. 2003. "What Difference Does the Choice of SES Make in Health Inequality Measurement?" *Journal of Health Economics* 12 (10): 885–890.
- Wall, S.N., and others. 2009. "Neonatal Resuscitation in Low-Resource Settings: What, Who, and How to Overcome Challenges to Scale Up?" International Journal of Gynecology and Obstetrics 107 (Suppl.): S47–S64.
- Wakabi, W. 2008. "Extension Workers Drive Ethiopia's Primary Health Care." *Lancet* 372 (9642): 880.
- WHO (World Health Organization). 1985. "Appropriate Technology for Birth." *Lancet* 326 (8452): 436–437.
- ——. 2005a. "Working Towards Universal Coverage of Maternal, Newborn and Child Health Interventions." World Health Assembly Resolution 58.31. World Health Organization, Geneva.
- . 2005b. World Health Report 2005: Make Every Mother and Child Count. Geneva: World Health Organization.
- ——. 2006. World Health Report 2006: Working for Health. Geneva: World Health Organization.

- ——. 2009. *Global Atlas of the Health Workforce.* Geneva: World Health Organization.
- —. 2010. *World Health Statistics 2010*. Geneva: World Health Organization.
- WHO (World Health Organization), Commission for Social Determinants of Health. 2008. "Social Determinants of Health." World Health Organization, Geneva. [http://www.who.int/social_determinants/en/].
- WHO (World Health Organization), Regional Office for South-East Asia. 2009. *Safer Pregnancy in Tamil Nadu: From Vision to Reality*. New Delhi: World Health Organization Regional Office for South-East Asia.
- WHO (World Health Organization) and UNICEF (United Nations Children's Fund). 2004a. "Clinical Management of Acute Diarrhoea." WHO–UNICEF joint statement. Geneva and New York: World Health Organization and United Nations Children's Fund.
- . 2004b. "Management of Pneumonia in Community Settings." WHO–UNICEF joint statement. Geneva and New York: World Health Organization and United Nations Children's Fund.
- . 2007. Report Card on the Prevention of Mother-to-ChildTransmission of HIV and Paediatric Care. Geneva and New York: World Health Organization and United Nations Children's Fund.
- . 2009. "Home Visits for the Newborn Child: A Strategy to Improve Survival." WHO–UNICEF joint

- statement. Geneva and New York: World Health Organization and United Nations Children's Fund.
- WHO (World Health Organization), UNFPA (United Nations Population Fund), UNICEF (United Nations Children's Fund), and AMDD (Averting Maternal Death and Disability). 2009. Indicators for Monitoring the Availability and Use of Obstetric Services: A Handbook. Geneva: World Health Organization.
- WHO (World Health Organization), UNICEF (United Nations Children's Fund), UNFPA (United Nations Population Fund), and World Bank. 2007. *Maternal Mortality in 2005*. Geneva: World Health Organization.
- WHO (World Health Organization–UNICEF (United Nations Children's Fund) Joint Programme on Water and Sanitation. 2010. *Progress on Sanitation and Drinking Water: 2010 Update.*Geneva and New York: World Health Organization and United Nations Children's Fund.
- Witter S., S. Adjei, M. Armar-Klemsesu, and W. Graham. 2009. "Providing Free Maternal Health Care: Ten Lessons from an Evaluation of the National Delivery Exemption Policy in Ghana." Global Health Action 2.
- Xu, K., D. Evans, G. Carrin, and A.M. Aguilar-Rivera. 2005. "Designing Health Financing Systems to Reduce Catastrophic Health Expenditures." Technical Brief for Policy Makers WHO/EIP/HSF/ PB/05.02. World Health Organization, Geneva.
- You D.,T. Wardlaw, P. Salama, and G. Jones. 2009. "Levels and Trends in Under-5 Mortality, 1990–2008." *Lancet* 375 (9709): 100–103.





